



## Prevalence of glaucoma in rural Myanmar: the Meiktila Eye Study

R J Casson, H S Newland, J Muecke, S McGovern, L Abraham, W K Shein, D Selva and T Aung

*Br. J. Ophthalmol.* 2007;91;710-714  
doi:10.1136/bjo.2006.107573

---

Updated information and services can be found at:  
<http://bjo.bmj.com/cgi/content/full/91/6/710>

---

*These include:*

### References

This article cites 28 articles, 15 of which can be accessed free at:  
<http://bjo.bmj.com/cgi/content/full/91/6/710#BIBL>

1 online articles that cite this article can be accessed at:  
<http://bjo.bmj.com/cgi/content/full/91/6/710#otherarticles>

### Rapid responses

You can respond to this article at:  
<http://bjo.bmj.com/cgi/eletter-submit/91/6/710>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Notes

---

To order reprints of this article go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *British Journal of Ophthalmology* go to:  
<http://journals.bmj.com/subscriptions/>

## WORLD VIEW

## Prevalence of glaucoma in rural Myanmar: the Meiktila Eye Study

R J Casson, H S Newland, J Muecke, S McGovern, L Abraham, W K Shein, D Selva, T Aung

*Br J Ophthalmol* 2007;**91**:710–714. doi: 10.1136/bjo.2006.107573

See end of article for authors' affiliations

Correspondence to:  
Professor R J Casson,  
Associate Professor,  
Department of  
Ophthalmology and Visual  
Sciences, South Australian  
Institute of Ophthalmology,  
Adelaide University,  
Adelaide 5000, South  
Australia, Australia; robert.  
casson@adelaide.edu.au

Accepted 1 November 2006

**Aim:** To determine the prevalence of glaucoma in the Meiktila district of central, rural Myanmar.**Methods:** A cross-sectional, population-based survey of inhabitants  $\geq 40$  years of age from villages in Meiktila district, Myanmar, was performed; 2481 eligible participants were identified and 2076 participated in the study. The ophthalmic examination included Snellen visual acuity, slit-lamp examination, tonometry, gonioscopy, dilated stereoscopic fundus examination and full-threshold perimetry. Glaucoma was classified into clinical subtypes and categorised into three levels according to diagnostic evidence.**Results:** Glaucoma was diagnosed in 1997 (80.5%) participants. The prevalence of glaucoma of any category in at least one eye was 4.9% (95% CI 4.1 to 5.7;  $n = 101$ ). The overall prevalence of primary angle-closure glaucoma (PACG) was 2.5% (95% CI 1.5 to 3.5) and of primary open-angle glaucoma (POAG) was 2.0% (95% CI 0.9 to 3.1). PACG accounted for 84% of all blindness due to glaucoma, with the majority due to acute angle-closure glaucoma (AACG).**Conclusion:** The prevalence of glaucoma in the population aged  $\geq 40$  years in rural, central Myanmar was 4.9%. The ratio of PACG to POAG was approximately 1.25:1. PACG has a high visual morbidity and AACG is visually devastating in this community. Screening programmes should be directed at PACG, and further study of the underlying mechanisms of PACG is needed in this population.

Glaucoma is the second most common cause of world blindness, and the majority of those blinded reside in Asia.<sup>1–2</sup> Recent studies have provided valuable information about the prevalence and subtypes of glaucoma in certain Asian regions,<sup>3–12</sup> and it has become recognised that angle-closure glaucoma is more common in people of Asian origin than those with European or African ethnicity<sup>5–13–16</sup>; however, the relative rates of open-angle to closed-angle glaucoma are region-dependent within Asia, with the rate of primary angle-closure glaucoma (PACG) particularly high in Mongolian and Chinese eyes,<sup>5–8–17</sup> and variable across India.<sup>6–10–12–18</sup> In accordance with the World Health Organization's (WHO's) Vision 20/20 initiative, the assessment of the prevalence of glaucoma subtypes is important because it has implications for the optimisation of screening programmes and treatment strategies.<sup>19–23</sup>

WHO estimates of the prevalence of glaucoma in many Asian regions are crude. Limited WHO data<sup>24</sup> and anecdotal evidence suggested high rates of angle-closure glaucoma in the Union of Myanmar (Myanmar; formerly Burma). Until now, no robust population-based data have been available on the prevalence and subtypes of glaucoma in Myanmar. Here, we report on the prevalence and subtypes of glaucoma in the inhabitants of the rural, central region of this country.

## METHODS

## Sampling procedure

The Meiktila Eye Study (MES) was a population-based, cross-sectional ophthalmic survey of the inhabitants of rural villages in central Myanmar. The principal aims of this project were to estimate the prevalence and causes of visual impairment, and the prevalence and risk factors of ocular disorders, including glaucoma, among persons  $\geq 40$  years of age in this region.

The study was conducted within the Mandalay Division, an area encompassing 34 253 km<sup>2</sup> divided into seven second-order administrative districts of approximately equal size. The township of Meiktila (population approximately 251 000), located at

20°53'N, 95°53' E, lies centrally in the Meiktila District, and is the only urban region in this entire district. The District is arbitrarily divided by the Ministry of Health (MOH) into six zones served by a centrally located eye hospital in Meiktila.

Participants were selected using a randomised, stratified, cluster sampling process. A sampling frame consisting of a list of all villages in the Meiktila District along with their populations was obtained from the MOH. Villages were arbitrarily stratified as large (population  $> 825$ ) or small (population  $\leq 825$ ), with small villages in each of the six zones within the Meiktila District constituting six separate strata. For logistical reasons, sampling was restricted to villages within 3 h drive from Meiktila (an area encompassing approximately 80% of the district). All persons aged  $\geq 40$  years from each selected village were eligible for inclusion. The sample size was based on the desired precision of the estimate of blindness (the principal aim of the MES); the assessment of glaucoma prevalence was a secondary objective. Healthcare workers from Meiktila township enumerated the selected villages (and advertised and promoted the survey) before commencement of the survey. Six small villages (one from each zone) and four large villages were enumerated, providing a total sample population of 2481 people.

## Data collection

Data collection was performed at the end of the rainy season in November 2005. A single survey team conducted the entire study. Each team member was assigned specific tasks and was well trained in the appropriate area. Specific observations were done by 1–2 members, limiting or eliminating interobserver

**Abbreviations:** AACG, acute angle-closure glaucoma; CDR, cup/disc ratio; FDT, frequency doubling technology; IOP, intraocular pressure; ISGEO, International Society for Geographic and Epidemiological Ophthalmology; MES, Meiktila Eye Study; MOH, Ministry of Health; PACG, primary angle-closure glaucoma; POAG, primary open-angle glaucoma; TM, trabecular meshwork; VA, visual acuity; WHO, World Health Organization

**Box 1 Diagnostic criteria for glaucoma**

- **Category 1 diagnosis** (structural and functional evidence): eyes with a cup:disc ratio (CDR) >97.5th centile for the normal (non-glaucomatous) population (CDR  $\geq 0.7$  was used on the basis of the data from previous studies in the region), or a CDR  $\geq 0.6$  in the presence of asymmetry  $\geq 0.3$  or a neuroretinal rim width reduced to <0.1 CDR (between 11:00 and 13:00 or between 17:00 and 19:00 h) and a definite visual field defect consistent with glaucoma. Eyes with evidence of previous acute angle-closure glaucoma (AACG) which had no perception of light (NPL) were also classified as category 1, even if the optic disc was not visualised ("end-stage" AACG).
- **Category 2 diagnosis** (advanced structural damage with unproved field loss): if the subject could not satisfactorily complete visual field testing, but had a CDR >99.5th centile for the normal (non-glaucomatous) population (CDR  $\geq 0.8$  was used on the basis of the data from the normal population in this study), glaucoma was diagnosed solely on the basis of structural evidence.
- **Category 3 diagnosis** (optic disc not seen; field test impossible): if it was not possible to examine the optic disc, glaucoma was diagnosed if: (A) the visual acuity (VA) was <3/60 and the intraocular pressure >99.5th centile; or (B) the VA was <3/60 and the eye showed evidence of glaucoma-filtering surgery.

variability. All equipment and personnel were transported to each village, and the data collection was performed on site. A medical and ophthalmic history was obtained from each patient in his or her own language by qualified healthcare workers. Each participant then received a comprehensive vision and eye examination.

Visual acuity (VA) was tested unaided, and with a pinhole using a well-illuminated Snellen chart at 6 m. Intraocular pressure (IOP) was measured with a Goldmann applanation tonometer (Haag-Streit, Koeniz, Switzerland) and anterior segment examination was performed using a slit lamp. The presence of previous iris ischaemia or pseudoexfoliation was recorded. Gonioscopy was performed by two experienced ophthalmologists using a Sussman gonioscope. Static gonioscopy was performed in dim illumination with minimal pressure on the cornea using a short slit beam; each quadrant was graded using the Scheie classification. If >90° of posterior trabecular meshwork (TM) was visible, the pupil was dilated with tropicamide 1% and phenylephrine 2.5%. Eyes with  $\leq 90^\circ$  of posterior TM visible were deemed "occludable" and dilated with tropicamide 0.5% only and kept under observation for 4 h;

**Table 1** Intraocular pressure in normal participants\*

	Right IOP (mm Hg) (95% CI)	Left IOP (mm Hg) (95% CI)
Number of measurements	1952	1953
Mean	14.8 (14.65 to 14.95)	14.9 (14.75 to 15.05)
Median	14	15
97.5th centile	21.7 (21.55 to 21.85)	21.9 (21.75 to 22.05)
99.5th centile	25.0 (24.85 to 25.15)	25.4 (25.25 to 25.55)

IOP, intraocular pressure.

\*Subjects had neither structural nor functional evidence of glaucomatous optic neuropathy.

if not possible, they were not dilated. If either eye had evidence of previous acute angle-closure glaucoma (AACG; see definition below), then neither eye was dilated. Optic disc and retinal examination was performed by two experienced ophthalmologists using a 78 D lens and reference to standard disc images. The vertical cup:disc ratio (CDR) and the presence of focal notching were recorded. The agreement between the two ophthalmologists was good for grading the occludability ( $\kappa = 0.78$  and determining the CDR ( $\kappa = 0.72$ ).

Eyes with VA >6/60, and which fulfilled category 1 optic disc criteria (see below), underwent full-threshold perimetry (C-20 strategy) using frequency doubling technology (FDT; Zeiss Humphrey Systems, Dublin, California, USA). Tests were considered reliable if there were <20% fixation errors and <33% false-positive and false-negative errors. All individuals were naïve to perimetry and received instruction in their own language, followed by a practice in the demonstration mode. If the initial test was unreliable, individuals were given a second attempt. More than one missed point on the pattern deviation was considered abnormal.<sup>25</sup>

**Ethics**

The MES was approved by the MOH in Myanmar and had ethical approval from the Royal Adelaide Hospital Ethics Committee. Consent for participation was obtained from the head of each village before commencement of the survey, and written, informed consent, in the participant's own language, was obtained from all willing participants. The study was conducted in accordance with the Declaration of Helsinki.

**Statistics**

Prevalence rates were calculated as ratio estimates using appropriate weights for each of the sampled villages. Bootstrapping was used to overcome the problem of variance estimation in clusters where only the one primary sampling unit (village) was selected. All prevalence estimates were calculated using SAS V.9.1. Villages were randomly selected; hence, point prevalences are unbiased.

**Definitions**

A three-tiered system of evidence, as suggested by the International Society for Geographic and Epidemiological Ophthalmology (ISGEO),<sup>26</sup> was used to categorise glaucoma (box 1).

Blindness due to glaucoma was defined as an eye with pinhole vision <3/60; fields were not taken into consideration.

Prevalence was calculated on an individual rather than a per eye basis. If at least one eye was diagnosable using the above criteria, the subject was included in the prevalence analysis. In those participants with only one diagnosable eye, if glaucoma was not present in this eye, they were assumed not to have glaucoma in the undiagnosable eye.

Glaucoma was also categorised into three principal clinical subtypes:

1. PACG was diagnosed if the criteria for category 1–3 were met,  $\leq 90^\circ$  of posterior TM was visible with static gonioscopy and no secondary cause for glaucoma was present. PACG was further subdivided into acute and chronic forms. Chronic PACG was diagnosed if the above criteria were met, and acute PACG was diagnosed if the above criteria were met and there was evidence of previous iris ischaemia (defined as the presence of iris whorling or stromal atrophy).<sup>27</sup> Historical evidence only of an attack of AACG was considered insufficient for diagnosis.
2. Primary open-angle glaucoma (POAG) was diagnosed if the criteria for category 1–3 were met, >90° of posterior TM

**Table 2** Vertical cup:disc ratio in normal participants\*

	Right CDR	Left CDR
Number of measurements	1850	1852
Mean	0.34	0.34
Median	0.3	0.3
97.5th centile	0.64	0.66
99.5th centile	0.79	0.82

CDR, cup:disc ratio.

\*Defined as those with available vertical CDR data and excluding eyes with definitive glaucomatous field defect.

was visible on static gonioscopy and no secondary cause for glaucoma was present.

- Secondary glaucoma was diagnosed if the criteria for category 1–3 were met and a secondary cause was evident. This included pseudoexfoliative and neovascular glaucoma.

## RESULTS

A total of 2481 participants were eligible and 2076 were examined (836 men, 1240 women; participation rate 83.7%). The mean age was 56.2 years. Sufficient examination data to diagnose glaucoma (as defined above) in at least one eye were obtained from 1997 participants. There were 110 participants in whom glaucoma was diagnosable in one eye only. The mean IOP and CDR for the normal (non-glaucomatous) population are shown in tables 1 and 2, respectively. There were seven eyes which had end-stage AACG and were classified in category 1.

The prevalence of glaucoma (allowing for the study design) in any category in at least one eye was 4.9% (95% CI 4.1 to 5.7; n = 101 participants). There were 156 eyes of 101 participants which met the ISGEO three-tiered evidence-based classification of glaucoma: 51 eyes were in category 1, 73 were in category 2 (perimetry not performed or unreliable) and 32 were in category 3. Of the 73 eyes in category 2, 46 did not meet the VA criteria (VA >6/60) and the remainder could not perform reliable (as defined above) perimetry by the second attempt. Seven eyes met the structural definitions of criteria 1, but did not meet the VA standard for perimetry or could not perform reliable perimetry, and were deemed non-glaucomatous. Only three eyes were classified as glaucomatous on the basis of CDR asymmetry.


The prevalence of glaucoma increased with age in both men and women (fig 1). Figure 2 shows the distribution of PACG, POAG and secondary glaucoma in men and women. The overall prevalence of PACG was 2.5% (95% CI 1.5 to 3.5) and of POAG was 2.0% (95% CI 0.9 to 3.1). In all, 22 (1.1%) participants had AACG in at least one eye and 30 (1.5%) had CACG. There were 10 (0.5%) participants with secondary glaucoma in at least one eye: 5 eyes with pseudoexfoliative, 3 with uveitic and 2 with neovascular glaucoma.

There were 32 eyes blinded by PACG, accounting for 84% of all eyes with blindness due to glaucoma. AACG was the cause of blindness in 20 eyes, and CACG in 12 eyes. Eight participants were bilaterally blind due to AACG and three due to CACG. Only three eyes were blind due to POAG and four due to secondary glaucoma.

The use of mydriatics, as per the protocol, produced no adverse events.


## DISCUSSION

This study provides the first population-based data about the prevalence and subtypes of glaucoma in Myanmar. Data relating to the prevalence of “occludable” angles (angle-closure glaucoma suspects) are not presented in this report. The most striking finding was the high prevalence of PACG in this

**Figure 1** Prevalence of glaucoma by age and gender.

population. The ratio of PACG to POAG was 1.25:1, which is lower than the 3:1 ratio reported in a Mongolian population,<sup>5</sup> but almost twice that reported in Chinese eyes,<sup>3, 8</sup> and considerably greater than the ratios reported in populations from India and Bangladesh (table 3).<sup>6, 7</sup> Discerning the relative amounts of PACG to POAG is important because it has profound implications in the optimisation of screening and treatment strategies: a greater prevalence of PACG, coupled with its high visual morbidity, implies that more resources should be directed towards it. A directed screening programme involving Van Herick grading, gonioscopy and laser iridotomy has been highly successful in reducing angle-closure glaucoma in the Inuit and is undergoing evaluation in Mongolia.<sup>19</sup> However, evidence is emerging which suggests that the mechanism of angle closure in certain regions of Asia, including South-East Asia, may be multifactorial, involving pupillary block and non-pupillary block components.<sup>19, 28</sup>

Although we recognise that consistency among epidemiological studies is important and have modelled this study, as much as practically possible, on similar studies from this region,<sup>3, 6-8</sup> we chose to slightly modify the ISGEO inclusion criteria: we included eyes with no perception of light (NPL; with evidence of old AACG) in which the optic disc was not seen, irrespective of the IOP, and which did not meet the criteria for classification in the current ISGEO system. In our opinion, these eyes have evidence of severe functional and implied structural optic neuropathy secondary to an old acute

**Figure 2** Distribution of clinical glaucoma subtypes by gender.

**Table 3** Prevalence of glaucoma in various Asian populations

Author (year)	Location	Number	Age group (years)	Prevalence of PACG	Prevalence of POAG	Ratio POAG:PACG
He <i>et al.</i> <sup>3</sup> 2006	Liwan, Guangzhou	1405	≥50	1.5%	2.1%	0.71:1
Vijaya <i>et al.</i> <sup>4, 10</sup> 2005, 2006	Chennai, India	3924	≥40	0.87%	1.62%	0.54:1
Raychaudhuri <i>et al.</i> <sup>6</sup> 2005	Calcutta, India	1269	≥50	0.72%	3.6%	0.2:1
Rahman <i>et al.</i> <sup>7</sup> 2004	Dhaka, Bangladesh	2346	≥35	0.4%	2.5%	0.16:1
Foster <i>et al.</i> <sup>8</sup> 2000	Tanjong Pagar, Singapore	1232	40–79	1.13%	1.78%	0.63:1
Dandona <i>et al.</i> <sup>11, 12</sup> 2000	Hyderabad, India	1399	≥40	1.08%	2.56%	0.41:1
Present study	Meiktila, Myanmar	1997	≥40	2.5%	2.0%	1.25:1

PACG, primary angle-closure glaucoma; POAG, primary open-angle glaucoma.

IOP increase and warrant classification in category 1. However, even if these eyes were excluded from the analysis, the overall prevalence (4.7%) is minimally affected.

The prevalence of glaucoma in this study is a little higher than prevalence rates reported in other population-based studies from Asia<sup>3–9</sup> (table 3). This may partly relate to the use of FDT perimetry, with relatively high sensitivity, and to the inclusion of eyes with old AACG. However, it may simply reflect a particularly high rate of PACG in this population.

Most of those eyes that were not adequately examined for glaucoma, had dense cataracts. Some of these eyes could also have had glaucoma so that the overall estimate for the prevalence of glaucoma (4.9%) may, in fact, be conservative. The 97.5th centile for the CDR in the normal population was approximately 6.5; however, based on recent data from this region a CDR of ≥0.7 was chosen as the cut-off for field testing; hence, early glaucoma with concentric cupping could have been missed. It is also likely that many of the participants classified as having PACG actually had combined-mechanism glaucoma; however, relationships between the amount of angle closure and the IOP, which may arouse suspicion of a combined mechanism, were not taken into consideration in this study.

The optimal method of perimetry for studies of this nature conducted “in the field” is unclear. We chose to use FDT because of its availability, portability, relative impunity to defocus,<sup>29</sup> ease of use and recent use in a similar population-based study in India.<sup>4, 10</sup> The current study was designed to detect glaucoma based on ISGEO criteria; hence, perimetry was only designated for those participants at least meeting the category 1 structural criteria. An arbitrary VA cut-off for perimetry was set at presenting Snellen acuity >6/60. Even at this level, almost 63% of eyes in category 2 were not eligible for perimetry, hence the relatively low rate (25%) of reliable perimetric data on the population meeting other diagnostic criteria. Previous similar studies<sup>4, 10</sup> using FDT had set the VA limit at 6/24; however, given that the FDT sensitivity suffers little from up to 6 D of defocus,<sup>29</sup> and the high prevalence of visual impairment in this population, our VA criterion seems reasonable. Arguably, the low rate of reliable perimetric data casts doubt on the prevalence of glaucoma; however, the ISGEO guidelines for the diagnosis of glaucoma are deliberately weighted towards structural changes, because it is well recognised that reliable perimetry in population-based studies, particularly in the developing world, is difficult.

Although the participation rate was relatively high (83%), we have no robust data about the visual status and ocular health of the non-participants. Anecdotally (according to the village chiefs), the principal reason for non-participation was occupation-related; hence, it is unlikely that any of the non-participants were glaucoma blind, suggesting that the prevalence of glaucoma in this group would be lower than in the participants. Although accurate data about the gender

distribution in the Meiktila district were not available, it is likely that women were over-represented in this study (59%), a common occurrence among similar studies, possibly reflecting occupation-related availability.

The Meiktila District was chosen for logistical reasons, not randomly, and may not be representative of neighbouring regions within the Mandalay Division of central Myanmar; however, we have no reason to believe that this is the case.

In conclusion, the prevalence of glaucoma in the population ≥40 years of age in the Meiktila District of rural, central Myanmar is 4.9%. The ratio of PACG to POAG is approximately 1.25:1. Given the high visual morbidity of PACG, screening programmes should be directed at this disease and further study of the underlying mechanisms of PACG in this population is needed.

#### ACKNOWLEDGEMENTS

We thank the Myanmar Ministry of Health for their invaluable assistance with this survey. This study was possible only due to the support of the staff at the Yangon Eye Hospital and Meiktila Eye Hospital. Expert assistance was provided by Drs Tun Aung Kyaw and Nyunt Maung from the Trachoma Control and Prevention of Blindness Program. Thanks to Paul Foster for generous and expert advice concerning the methodology and technical aspects. Australian Ambassador to Yangon, Bob Davis, provided valuable assistance and donated a slit lamp to the Meiktila Eye Hospital. We thank Alcon Australia for the loan of equipment. The sole financial support was provided by a generous grant from Pfizer Ophthalmic. We thank the people who participated in this study and who welcomed us so kindly into their villages.

#### Authors' affiliations

**R J Casson, H S Newland, J Muecke, S McGovern, I Abraham, D Selva**, Department of Ophthalmology and Visual Sciences, South Australian Institute of Ophthalmology, Adelaide University, Adelaide, South Australia, Australia

**W K Shein**, Meiktila Eye Hospital and Trachoma Control and Prevention Programme, Meiktila, Myanmar

**T Aung**, Yangon Eye Hospital, Yangon, Myanmar

Funding: The survey was funded by a grant from Pfizer Ophthalmic. The design of the survey, its execution, analysis, interpretation and publication were carried out independently by the authors.

Competing interests: None.

#### REFERENCES

- 1 Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol* 2006;**90**:262–7.
- 2 Resnikoff S, Pascolini D, Etya'ale D, *et al.* Global data on visual impairment in the year 2002. *Bull World Health Organ* 2004;**82**:844–51.
- 3 He M, Foster PJ, Ge J, *et al.* Prevalence and clinical characteristics of glaucoma in adult Chinese: a population-based study in Liwan district, Guangzhou. *Invest Ophthalmol Vis Sci* 2006;**47**:2782–8.
- 4 Vijaya L, George R, Arvind H, *et al.* Prevalence of angle-closure disease in a rural southern Indian population. *Arch Ophthalmol* 2006;**124**:403–9.

- 5 Foster PJ, Baasanhu J, Alsbirk PH, *et al*. Glaucoma in Mongolia. A population-based survey in Hovsgol province, northern Mongolia. *Arch Ophthalmol* 1996;**114**:1235-41.
- 6 Raychaudhuri A, Lahiri SK, Bandyopadhyay M, *et al*. A population based survey of the prevalence and types of glaucoma in rural West Bengal: the West Bengal Glaucoma Study. *Br J Ophthalmol* 2005;**89**:1559-64.
- 7 Rahman MM, Rahman N, Foster PJ, *et al*. The prevalence of glaucoma in Bangladesh: a population based survey in Dhaka division. *Br J Ophthalmol* 2004;**88**:1493-7.
- 8 Foster PJ, Oen FT, Machin D, *et al*. The prevalence of glaucoma in Chinese residents of Singapore: a cross-sectional population survey of the Tanjong Pagar district. *Arch Ophthalmol* 2000;**118**:1105-11.
- 9 Bourne RR, Sukudom P, Foster PJ, *et al*. Prevalence of glaucoma in Thailand: a population based survey in Rom Klao District, Bangkok. *Br J Ophthalmol* 2003;**87**:1069-74.
- 10 Vijaya L, George R, Paul PG, *et al*. Prevalence of open-angle glaucoma in a rural south Indian population. *Invest Ophthalmol Vis Sci* 2005;**46**:4461-7.
- 11 Dandona L, Dandona R, Mandal P, *et al*. Angle-closure glaucoma in an urban population in southern India. The Andhra Pradesh Eye Disease Study. *Ophthalmology* 2000;**107**:1710-16.
- 12 Dandona L, Dandona R, Srinivas M, *et al*. Open-angle glaucoma in an urban population in southern India: the Andhra Pradesh Eye Disease Study. *Ophthalmology* 2000;**107**:1702-9.
- 13 Salmon JF, Mermoud A, Ivey A, *et al*. The prevalence of primary angle closure glaucoma and open angle glaucoma in Mamre, western Cape, South Africa. *Arch Ophthalmol* 1993;**111**:1263-9.
- 14 Foster PJ. The epidemiology of primary angle closure and associated glaucomatous optic neuropathy. *Semin Ophthalmol* 2002;**17**:50-8.
- 15 Bourne RR, Sorensen KE, Klauber A, *et al*. Glaucoma in East Greenlandic Inuit—a population survey in Iltoqqortoormiit (Scoresbysund). *Acta Ophthalmol Scand* 2001;**79**:462-7.
- 16 Chew PT, Aung T. Primary angle-closure glaucoma in Asia. *J Glaucoma* 2001;**10**:57-8.
- 17 Foster PJ, Johnson GJ. Glaucoma in China: how big is the problem? *Br J Ophthalmol* 2001;**85**:1277-82.
- 18 Thulasiraj RD, Nirmalan PK, Ramakrishnan R, *et al*. Blindness and vision impairment in a rural south Indian population: the Aravind Comprehensive Eye Survey. *Ophthalmology* 2003;**110**:1491-8.
- 19 Johnson GJ, Foster PJ. Can we prevent angle-closure glaucoma? *Eye* 2005;**19**:1119-24.
- 20 Aung T, Chew PT. Review of recent advancements in the understanding of primary angle-closure glaucoma. *Curr Opin Ophthalmol* 2002;**13**:89-93.
- 21 Nolan WP, Aung T, Machin D, *et al*. Detection of narrow angles and established angle closure in Chinese residents of Singapore: potential screening tests. *Am J Ophthalmol* 2006;**141**:896-901.
- 22 Friedman DS, Gazzard G, Foster P, *et al*. Ultrasonographic biomicroscopy, Scheimpflug photography, and novel provocative tests in contralateral eyes of Chinese patients initially seen with acute angle closure. *Arch Ophthalmol* 2003;**121**:633-42.
- 23 Saw SM, Gazzard G, Friedman DS. Interventions for angle-closure glaucoma: an evidence-based update. *Ophthalmology* 2003;**110**:1869-78.
- 24 Alsbirk PH. *Prevention and control of visual impairment and blindness (with special reference to glaucoma) in Burma, Consultant Report*. World Health Organization, Southeast Asia Region/Ophthalmology, 1984.
- 25 Muskens RP, Heeg GP, Jansonius NM. An evaluation of algorithms designed to classify the results from frequency doubling perimetry. *Ophthalmic Physiol Opt* 2004;**24**:498-503.
- 26 Foster PJ, Buhmann R, Quigley HA, *et al*. The definition and classification of glaucoma in prevalence surveys. *Br J Ophthalmol* 2002;**86**:238-42.
- 27 Loon SC, Chew PT, Oen FT, *et al*. Iris ischaemic changes and visual outcome after acute primary angle closure. *Clin Experiment Ophthalmol* 2005;**33**:473-7.
- 28 Aung T, Nolan WP, Machin D, *et al*. Anterior chamber depth and the risk of primary angle closure in 2 East Asian populations. *Arch Ophthalmol* 2005;**123**:527-32.
- 29 Anderson AJ, Johnson CA. Frequency-doubling technology perimetry and optical defocus. *Invest Ophthalmol Vis Sci* 2003;**44**:4147-52.

## VIDEO REPORT .....

*Br J Ophthalmol* 2007;**91**:714. doi: 10.1136/bjo.2007.121343

### Descemet-Stripping Automated Endothelial Keratoplasty Technique in Patients with Anterior Chamber Intraocular Lenses

Brian Groat, Michelle S Ying, David T Vroman, Luis E Fernández de Castro  
Storm Eye Institute, Medical University of South Carolina, Charleston, SC, USA

#### ABSTRACT

**Background:** Descemet-Stripping Automated Endothelial Keratoplasty (DSAEK) is a relatively new, but proven procedure in treating endothelial dysfunction. Patients with aphakia or anterior chamber intraocular lenses (ACIOL) pose more of a challenge when performing DSAEK. We report a technique that can be used during DSAEK to address situations.

**Case Report:** An 80-year-old white female with a history of intracapsular cataract extraction and a secondary ACIOL in her left eye was referred with pseudophakic bullous keratopathy and 3+ stromal edema. DSAEK was performed with two modifications: 1) occlusive pupilloplasty, and 2) a fixation suture to stabilize the donor tissue during the unfolding process. Her two-week postoperative visual acuity was 20/60.

**Video:** The surgical video demonstrates the modified DSAEK technique. A 10-0 polypropylene suture was used to create the occlusive pupilloplasty with a slipknot. The endothelium and Descemet were stripped using a 90-degree scraper. The donor tissue was folded in a 60/40 taco fashion and inserted into the anterior chamber. To stabilize the graft while unfolding in a shallow anterior chamber, a fixation suture was placed in the inferior portion of the graft. Air was injected, and the tissue unfolded in good position with the Lindstrom roller, and finally the occlusive pupilloplasty was released.

**Comment:** Currently, some patients with ACIOL are managed by performing an intraocular lens exchange with a scleral-sutured lens followed 6-8 weeks later by DSAEK. However, the two modifications described above allows DSAEK to be performed successfully in a single procedure with minimal additional surgical time.

To view the full report and accompanying video please go to: <http://bjo.bmj.com/cgi/content/full/91/6/714/DC1>

All videos from the **BJO video report collection** are available from: <http://bjo.bmj.com/video/collection.dtl>