

ACCEPTED VERSION

Haifeng Li, Wanqi Liang, Yun Hu, Lu Zhu, Changsong Yin, Jie Xu, Ludovico Dreni, Martin M. Kater, and Dabing Zhang

Rice MADS6 interacts with the floral homeotic genes SUPERWOMAN1, MADS3, MADS58, MADS13, and DROOPING LEAF in specifying floral organ identities and meristem fate

The Plant Cell, 2011; 23(7):2536-2552

© 2011 American Society of Plant Biologists. All rights reserved.

DOI: <http://dx.doi.org/10.1105/tpc.111.087262>

PERMISSIONS

http://tpc.msubmit.net/cgi-bin/main.plex?form_type=display_auth_instructions

ASPB is now offering an OPEN option that allows authors to have their articles available for free to all users immediately upon publication.

Authors are able to make the final publisher's version of the article free upon publication via our standard fee. Please note that ASPB retains copyright. Therefore, authors seeking to comply with institutional and other funder (e.g., RCUK) policies may deposit their peer-reviewed accepted manuscript into the appropriate repository for immediate access. Authors who are required to retain copyright and grant to their institution a nonexclusive license to exercise any and all rights under copyright, on a commercial or noncommercial basis, must file for a waiver of the institutional license.

11 June 2015

<http://hdl.handle.net/2440/92120>

Rice *MADS6* Interacts with the Floral Homeotic Genes *SUPERWOMAN1*, *MADS3*, *MADS58*, *MADS13*, and *DROOPING LEAF* in Specifying Floral Organ Identities and Meristem Fate

Haifeng Li,^{a,1} Wanqi Liang,^{a,1} Yun Hu,^a Lu Zhu,^a Changsong Yin,^a Jie Xu,^a Ludovico Dreni,^b Martin M. Kater,^b and Dabing Zhang^{a,c,2}

^aInstitute of Plant Science, State Key Laboratory of Hybrid Rice, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

^bDipartimento di Scienze Biomolecolari e Biotecnologie, Università degli Studi di Milano, 20133 Milan, Italy

^cBio-X Center, Key Laboratory of Genetics and Development and Neuropsychiatric Diseases, Ministry of Education, Shanghai Jiao Tong University, Shanghai 200240, China

AGAMOUS-LIKE6 (*AGL6*) genes play essential roles in flower development, but whether and how they work with floral organ identity genes remain less understood. Here, we describe interactions of the rice (*Oryza sativa*) *AGL6* gene *MADS6* with other rice floral homeotic genes in flower development. Genetic analyses revealed that *MADS6* specifies the identity of the three inner whorls and floral meristem determinacy redundantly with *SUPERWOMAN1/MADS16* (B-gene) or *MADS3* (C-gene). *MADS6* was shown to define carpel/ovule development and floral determinacy by interacting with *MADS13* (D-gene) and control the palea and floral meristem identities together with the *YABBY* gene *DROOPING LEAF*. Expression analyses revealed that the transcript levels of six B-, C-, and E-class genes were reduced in *mads6-1* at the early flower developmental stage, suggesting that *MADS6* is a key regulator of early flower development. Moreover, *MADS6* can directly bind to a putative regulatory motif on *MADS58* (C-gene), and *mads6-1 mads58* displayed phenotypes similar to that of *mads6-1*. These results suggest that *MADS6* is a key player in specifying flower development via interacting with other floral homeotic genes in rice, thus providing new insights into the mechanism by which flower development is controlled.

INTRODUCTION

The flowers of angiosperms display a huge diversity in structure (Theissen and Melzer, 2007). The ABCDE model proposed that the combinatorial action of class A, B, C, and E floral homeotic genes determines floral organ identity in eudicot model species, such as *Arabidopsis thaliana*, *Petunia hybrida*, and *Antirrhinum majus*. In particular, A- and E-function genes specify the identity of sepals in the first outer whorl; A-, B-, and E-function genes together determine the identity of petals in the second whorl; B-, C-, and E-function genes coordinately define stamen identity in the third whorl; and C- and E-function genes act together to specify carpels in the fourth whorl (Coen and Meyerowitz, 1991; Pelaz et al., 2000; Theissen, 2001; Theissen and Saedler, 2001; Ditta et al., 2004).


As one of the largest families in higher plants, the grass family (Poaceae) contains many economically important crops, such as rice (*Oryza sativa*), barley (*Hordeum vulgare*), and maize (*Zea mays*) (Clayton and Renvoize, 1986; Linder and Rudall, 2005). These crops evolved floral organization and morphologies distinct from those of eudicots and even other monocots (Grass Phylogeny Working Group, 2001; Rudall et al., 2005; Whipple et al., 2007). Spikelet, the structural unit of grass flowers, has a varied number of bract-like organs and florets. In rice, a spikelet consists of two pairs of sterile glumes (i.e., rudimentary glumes and empty glumes) and one floret that contains a lemma and a palea in whorl 1, two lodicules in whorl 2, six stamens in whorl 3, and a pistil in whorl 4 (Yuan et al., 2009; Zhang and Wilson, 2009). However, the molecular basis underlying grass flower development remains poorly understood (Clifford, 1987; Whipple et al., 2007).


Recently, emerging evidence suggests that the ABCDE genetic model is partially applicable to grasses such as rice and maize (Kozuka et al., 2000; Kater et al., 2006; Preston and Kellogg, 2006; Prusinkiewicz et al., 2007; Reinheimer and Kellogg, 2009; Thompson and Hake, 2009; Zhang and Wilson, 2009). For instance, mutations in maize *Silky1* and rice *SUPERWOMAN1* (*SPW1*) or *MADS16*, both of which are orthologs of the *Arabidopsis* B-function gene, *AP3*, cause homeotic transformations of stamens to carpels and lodicules to lemma- or palea-like structures (Ambrose et al., 2000; Nagasawa et al., 2003; Whipple et al., 2004), suggesting the conserved role of the B-class genes


¹These authors contributed equally to this work.

²Address correspondence to zhangdb@sjtu.edu.cn.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Dabing Zhang (zhangdb@sjtu.edu.cn).

Some figures in this article are displayed in color online but in black and white in the print edition.

Online version contains Web-only data.

Open Access articles can be viewed online without a subscription. www.plantcell.org/cgi/doi/10.1105/tpc.111.087262

from grasses to *Arabidopsis*. In grasses, there are duplicated and subfunctionalized C-class genes (Kramer et al., 2004; Zahn et al., 2006). For example, rice contains two AG homologs, *MADS3* and *MADS58* (Kramer et al., 2004). *MADS3* plays a key role in stamen and ovule identity specification, late anther development, and floral meristem determinacy (Yamaguchi et al., 2006; Hu et al., 2011; Li et al., 2011). Using an RNA interference approach, *MADS58* was shown to be required for defining floral meristem determinacy and carpel architecture (Yamaguchi et al., 2006). Similarly, maize has three AG homologs: *zag1* (*zea agamous1*), *zmm2* (*Zea mays mads2*), and *zmm23* (Münster et al., 2002). The *zag1* gene is required for floral meristem determinacy, but the biological functions of *zmm2* and *zmm23* have not been elucidated (Mena et al., 1996). Rice contains two D-class genes, *MADS13* and *MADS21*, which are orthologous to the *Arabidopsis* *SEEDSTICK* (*STK*) and petunia *FLORAL BINDING PROTEIN7* (*FBP7*) and *FBP11* genes (Colombo et al., 1995). *MADS13* was shown to be involved in ovule identity specification and floral meristem termination (Dreni et al., 2007; Li et al., 2011). However, mutants of the *Arabidopsis* *STK* gene do not display altered ovule identity (Pinyopich et al., 2003).

Grasses have diversified E-class (*SEPALLATA* [*SEP*]) genes. Rice has at least five *SEP*-like genes: *MADS1/LEAFY HULL STERILE1* (*LHS1*), *MADS5*, *MADS7*, *MADS8*, and *MADS34* (Malcomber and Kellogg, 2004, 2005; Zahn et al., 2005; Arora et al., 2007). *LHS1* specifies the identity of lemma and palea and the meristem of inner floral organs (Jeon et al., 2000; Agrawal et al., 2005; Prasad et al., 2005; Chen et al., 2006a). Transgenic plants with reduced expression of both *MADS7* and *MADS8* exhibit late flowering, homeotic transformations of lodicules, stamens, and carpels into palea/lemma-like organs, and a loss of floral determinacy. Simultaneous reduction of the expression of four rice *SEP*-like genes (i.e., *LHS1*, *MADS5*, *MADS7*, and *MADS8*) causes homeotic transformation of all floral organs except the lemma into leaf-like organs (Cui et al., 2010). *MADS34* (*PANICLE PHYTOMER2*) controls the development of inflorescences and spikelets (Gao et al., 2010; Kobayashi et al., 2010). Analysis of *mads34 mads1* indicates that *MADS34* and *LHS1* redundantly specify the identities of floral organs, including lemma/palea, lodicules, stamens, and carpel (Gao et al., 2010).

Sequence and phylogenetic analyses indicated that *AGAMOUS-LIKE6* (*AGL6*)-like and *SEP*-like genes have high sequence similarities, forming sister clades on the phylogenetic tree (Theissen et al., 2000; Becker and Theissen, 2003; Zahn et al., 2005). *SEP*-like genes are only found in angiosperms, while *AGL6* genes are ancient and widely distributed in gymnosperms and angiosperms. Recently, *AGL6*-like genes in monocots and eudicots were shown to play essential roles in flower development (Schauer et al., 2007; Ohmori et al., 2009; Reinheimer and Kellogg, 2009; Rijpkema et al., 2009; Li et al., 2010). The *Arabidopsis* genome contains two *AGL6* genes, *AGL6* and *AGL13* (Vandenbussche et al., 2003a), suggesting possible functional redundancy between the two genes. Mutation or knockdown of *AGL6* or *AGL13* does not result in an abnormal flower phenotype (Schauer et al., 2007; Koo et al., 2010; Yoo et al., 2011). Loss-of-function mutants of the only *Petunia* *AGL6* gene, Ph *AGL6*, show no morphological abnormalities of floral organs, but Ph *AGL6* functions redundantly with the *SEP* genes *FBP2* and *FBP5* in petal and

anther development, and its protein physically interacts with *FBP2* (Vandenbussche et al., 2003b; Rijpkema et al., 2009).

The *AGL6*-like genes from grasses form two paralogous clades: the *MADS17* clade containing only the rice *MADS17* gene, and the *MADS6* clade, which includes rice *MADS6* (also called *MOSAIC FLORAL ORGANS1* [*MFO1*]) and maize *ZAG3* and *ZAG5* (Ohmori et al., 2009; Reinheimer and Kellogg, 2009; Li et al., 2010). Grass *AGL6*-like genes were shown to be expressed in paleas, lodicules, ovules, and floral meristems, and each of these expression domains may represent a distinct function of the gene product (Reinheimer and Kellogg, 2009). In rice, the expression level of *MADS6* is high in floral meristem at early stages and in the palea and inner floral organ primordia (lodicule, stamen, and pistil) at later stages (Ohmori et al., 2009; Reinheimer and Kellogg, 2009; Li et al., 2010). *MADS17* transcripts were detected in the floral meristem at the early stage and in the lemma, palea, lodicule, pistil, and (weakly) in empty glumes and stamens at late stages, with its protein product functioning redundantly with *MADS6* in flower development (Ohmori et al., 2009; Reinheimer and Kellogg, 2009). Our previous studies revealed that the palea of *mads6-1* flowers develops five to six vascular bundles, which resembles the identity of a wild-type lemma, suggesting the role of *MADS6* in specifying the identity of palea. In addition, *mads6-1* flowers are retarded in development at the early stage, exhibit homeotic conversion of lodicules and stamens into glume-like and mosaic structures, have defective carpels and ovules, and contain indeterminate meristem at later flower developmental stages. Furthermore, we showed that the *MADS6* gene is able to specify floral state by determining floral organ and meristem identities together with *LHS1/MADS1* because *mads1-z mads6-1* double mutants display severe floral defects, such as no inner floral organs or glume-like structures within flowers and strongly indeterminate floral meristem, phenotypes not observed in the single mutants (Li et al., 2010). A mutation of the maize *AGL6* gene *zea agamous3* (*zag3*) results in extra mosaic or fused floral organs in the upper floral meristem and additional floral meristems in the lower floral meristem. *zag3* and the maize homolog of AG, *zag1*, can genetically and physically interact in promoting floral meristem identity (Thompson et al., 2009). These findings suggest that *AGL6* genes have *SEP*-like functions in flower development.

Despite the findings that *AGL6*-like genes have a role in defining floral organ and meristem identities, whether and how they interact with other floral homeotic genes in these processes remain largely unknown. Here, we report that *MADS6* interacts with several known flower homeotic genes in specifying flower development and determining floral meristem fate in rice. We show that *MADS6* not only interacts with B-, D-, and E-class proteins but also regulates the expression of these genes, thus providing novel insights into the mechanism by which *AGL6*-like genes exert their functions in plant flower development.

RESULTS

Transcriptome Analysis of *mads6-1* Flowers

To further elucidate the regulatory role of *MADS6*, we compared genome-wide mRNA levels in wild-type and *mads6-1* flowers

at stage Sp6, when stamen primordia are formed, using microarray analyses with an Agilent 4×4 4K oligonucleotide DNA chip. Stage Sp6 flowers were collected according to spikelet length and flower morphology defined by Ikeda et al. (2004), and three independent biological replicates were performed to assess its reproducibility. Data were analyzed by the Empirical Bayes method (Smyth, 2004). Initial filtering of candidate genes was performed using a false discovery rate cutoff of 0.5%, followed by a secondary selection using at least twofold changes in gene expression as the cutoff. Fifty-nine genes were found to have at least twofold changes in expression in *mads6-1* flowers compared with the wild type. Among them, 26 were upregulated and 33 were downregulated (see Supplemental Table 1 online).

The expression of six MADS box genes, including the B-class genes *MADS4* and *SPW1*, C-class genes *MADS3* and *MADS58*, and E-class genes *MADS7* and *MADS8*, was downregulated in the mutant (Table 1). In particular, *MADS58* and *MADS7* were downregulated ~22.5-fold (for one *MADS58* probe of chip) and 11.6-fold, respectively. The microarray data were further confirmed by serial quantitative RT-PCR (qRT-PCR) analysis, which showed that the mRNA levels of these six genes were reduced from stage 4 to stage 6 and came back up at stage 8 (Figure 1).

To clarify the functional relationship between *MADS6* and the floral homeotic genes whose expressions were significantly downregulated in *mads6-1*, we conducted detailed genetic analyses using double mutants between *mads6-1* and *spw-1*, *mads3-4*, *mads58*, *mads13-3*, and *dl-sup6*. In addition to phenotypic analysis, we also performed in situ analysis to determine the regulatory relationship between *MADS6* and *SPW1*, *MADS3*, *MADS58*, *MADS13*, and *DL* at the transcription level.

Interaction between *MADS6* and *SPW1*

SPW1 is a B-class gene required for determining the identity of lodicules and stamens. In *spw1-1* flowers, lodicules are transformed into glume-like structures and stamens into carpel-like organs (Nagasawa et al., 2003) (Figures 2A, 2D, 2P, and 2R). *mads6-1 spw1-1* flowers displayed enlarged palea and glume-like organs, which are similar to the phenotypes of *mads6-1* at stage Sp8 (Figures 2A to 2C, 2E, 2Q, and 2S) (Ohmori et al., 2009; Li et al., 2010), suggesting that *MADS6* specifies palea identity, and this function is independent from *SPW1*. Unlike *spw1-1* or *mads6-1*, in which the lodicule identity is partially lost, no lodicule-like organs were observed in whorl 2 of *mads6-1*

Table 1. Microarray Data Showing the Downregulated Expression of Six MADS Box Genes at Stage Sp6 in *mads6-1* Flower

Gene Name	Fold Change
<i>MADS16</i>	-8.4
<i>MADS4</i>	-3.7
<i>MADS3</i>	-7.7
<i>MADS58</i>	-22.5 ^a
<i>MADS58</i>	-6.1 ^a
<i>MADS7</i>	-11.6
<i>MADS8</i>	-6.3

^aDifferent probes for *MADS58*.




Figure 1. Expression Analysis of Six MADS Box Genes in *mads6-1* Flowers.

qRT-PCR results showing the relative expression levels of *SPW1*, *MADS4*, *MADS3*, *MADS58*, *MADS7*, and *MADS8* in *mads6-1* flowers at stage 4/5, stage 6, early stage 8, and late stage 8. The expression level of each gene in the wild type control was set as 100%, and error bars indicate SD ($n = 3$).

spw1-1 flowers (Figure 2F; see Supplemental Table 2 online), suggesting that *MADS6* and *SPW1* synergistically specify lodicule identity. Similar to *spw1-1*, *mads6-1 spw1-1* flowers displayed the conversion of stamens into carpel-like structures, each of which contains one to four stigmas (Figures 2F and 2S). Consistent with this phenotype, the expression of the carpel marker gene *DL* was found in the ectopic organs (see Supplemental Figures 1K to 1M online). These results suggest that *SPW1* plays a more important role in stamen identity. In the double mutant, the average number of glume-like organs per flower was 5.86 ($n = 43$) compared with 2.09 in *spw1-1* and 4.24 in *mads6-1* (see Supplemental Table 2 online). The average number of carpel-like organs per flower was 5.07 ($n = 43$) in the double mutant, while it was 7.00 for *spw1-1* ($n = 43$) and 1.10 for *mads6-1* ($n = 43$) (see Supplemental Table 2 online). More interestingly, about half of the flowers (22 in 43) in the double mutant developed new inflorescence-like organs at the position of wild-type lodicules (Figures 2G and 2H), a phenotype that was not observed in single mutant flowers, implying that *MADS6* and *SPW1* can repress the inflorescence primordia in the second whorl in a redundant manner. These observations suggested that both *MADS6* and *SPW1* function in specifying the identity of floral organs in the three inner whorls and the determinacy of floral meristem. Furthermore, the functions of these two proteins are redundant in some while independent from each other in other aspects of flower development.

To analyze phenotypes of the double mutant in more depth, we performed scanning electron microscopy. No structural differences between the flowers of *spw1-1* and the wild type were detected at stage Sp6, when stamen primordia initiate (Figure 2I). However, similar to that of *mads6-1* (Li et al., 2010), the inner flower organ primordia of the double mutant displayed retarded growth (Figure 2K). At stage Sp8, *spw1-1* displayed ectopic




Figure 2. Phenotypes of *spw1-1* and *mads6-1 spw1-1* Flowers.

(A) Wild-type (WT) flower with one lemma, one palea, two lodicules, six stamens, and one pistil.

(B) *mads6-1* flower with abnormal palea.

(C) and (D) Typical flower of *mads6-1* (C) and *spw1-1* (D) in which the lemma and the palea were removed to show the inner organs.

(E) *mads6-1 spw1-1* flower.

(F) and (G) *mads6-1 spw1-1* flowers of the type I (49%, $n = 43$) (F) and type II phenotype (51%, $n = 43$) (G). Lemma and palea were removed.

(H) to (O) Scanning electron microscopy analysis.

(H) A *mads6-1 spw1-1* flower showing the new inflorescence formed on the lemma side.

(I) and (J) *spw1-1* flowers at stage Sp6 (I) and Sp8 (J).

(K) and (L) *mads6-1 spw1-1* flower at stage Sp6 (K) and early stage of Sp8 (L).

(M) to (O) *mads6-1 spw1-1* flowers showing the initiation and development of additional inflorescence-like primordia at the position of lodicule.

(P) and (Q) Diagrams of wild-type (P) and *mads6-1* (Q) flowers.

(R) and (S) Diagrams of *spw1-1* (R) and *mads6-1 spw1-1* (S) flowers.

apa, abnormal palea; ca, carpel; ec, ectopic carpel; gl, glume-like organ; le, lemma; lo, lodicules; l-a, lodicule-anther mosaic organ; pl, palea-like organ; si, secondary inflorescence; sip, secondary inflorescence primordium; sp, spikelet; st, stamen. Bars = 1 mm in (A) to (G), 100 μm in (H), (J), and (L) to (O), 50 μm in (I) and (K).

carpel primordia (Figure 2J), and the double mutant flowers developed into carpel- and glume-like structures (Figures 2L and 2M). Also, additional inflorescence primordia were detectable in some *mads6-1 spw1-1* flowers (Figures 2N and 2O). Results of the scanning electron microscopy experiment were consistent with the genetic analysis, further substantiating the role *MADS6* and *SPW1* in regulating flower development.

RNA in situ hybridization analysis of the *mads6-1* mutant revealed that at stage Sp5, when lodicule primordia are formed, the expression level of *SPW1* in the lodicule primodium was slightly weaker than in the wild type (Figures 3A and 3E) (Nagasawa et al., 2003). At stage Sp6, the expression of *SPW1* was detected in lodicule and stamen primordia of *mads6-1* (Figures 3B and 3F). At stages Sp7 and Sp8, the expression of *SPW1* was observed in lodicule and stamen primordia in *mads6-1* similar with its expression pattern in the wild type (Figures 3C, 3D, 3G, and 3H). In *spw1-1*, the expression of *MADS6* was detected in paleas, lodicules, carpels, and the receptacle similar to its

expression profile in the wild-type plants (Figures 3I to 3L) (Ohmori et al., 2009; Reinheimer and Kellogg, 2009; Li et al., 2010). These in situ analysis results are consistent with the microarray and qRT-PCR experiments, supporting the notion that *MADS6* is able to activate the expression of *SPW1* at the early flower developmental stage but does not obviously affect its expression at late stages.

***MADS6* and *DL* Redundantly Regulate Floral Meristem Identity**

DL, which belongs to the *YABBY* gene family, determines carpel identity and regulates vascular pattern of the lemma (Nagasawa et al., 2003; Yamaguchi et al., 2004; Li et al., 2011). In the construction of double mutants, we used the null allele *dl-sup6*, which is allelic to the previously reported *dl-2* mutant (Nagasawa et al., 2003; Yamaguchi et al., 2004; Li et al., 2011). *dl-sup6* has drooping leaves, ectopic stamens in the fourth




Figure 3. Expression Pattern of *SPW1/MADS16* and *MADS6*.

(A) to (D) *SPW1* expression in wild-type (WT) flowers at stage Sp5 (A), Sp6 (B), Sp7 (C), and Sp8 (D).

(E) to (H) Expression of *SPW1* in *mads6-1* flowers at Sp5 (E), Sp6 (F), Sp7 (G), and Sp8 (H).

(I) to (L) Expression pattern of *MADS6* in *spw1-1* flowers at stage Sp5 (I), Sp6 (J), Sp7 (K), and Sp8 (L).

ca, carpel; ec, ectopic carpel; gl, glume-like organ; le, lemma; ll, lodicule-like organ; lo, lodicules; pl, palea-like organ; pa, palea; si, secondary inflorescence; sip, secondary inflorescence primordium; sp, spikelet; st, stamen. Bars = 50 μ m in (A), (E), and (I) and 100 μ m in (B) to (D), (F) to (H), and (J) to (L). Numbers in (J) to (L) indicate the whorl number.

whorl at the position of the carpel, and loss of floral meristem determinacy in some of the flowers (Figures 4A, 4B, and 4J) (Li et al., 2011).

Flowers of the *mads6-1 dl-sup6* double mutant could be divided into two types based on palea morphology. Type I was found in 58% of the flowers ($n = 56$). They displayed widened palea, which is similar to that of *mads6-1* (Figure 4C). Type II flowers (42%; $n = 56$) had two palea-like organs (Figures 4D and 4K), which were not observed in the single mutants. Examination of transverse sections showed that these two palea-like organs had the characteristic marginal tissue of the lemma but contained three vascular bundles similar to that of wild-type palea (Figures 4I and 4K). Consistent with this finding, two palea-like primordia were observed at stage Sp6 on the palea side in type II flowers (Figure 4G). These results revealed that *MADS6* specifies the palea identity together with *DL*.

The phenotypes of the *mads6-1 dl-sup6* floral organs in whorls 2 and 3 appeared similar to that of *mads6-1* (Figure 4E), suggesting that *MADS6* is involved in defining the identities of lodicules and stamens, whereas *DL* does not play a role in it. This

finding is consistent with the lack of *DL* expression in whorls 2 and 3 (see Supplemental Figure 1 online). Interestingly, all flowers in *mads6-1 dl-sup6* displayed an inflorescence-like structure in whorl 4 ($n = 62$) (Figures 4E, 4F, 4H, and 4K), yet this phenotype was rarely observed in *mads6-1* (Li et al., 2010). These results suggested that *MADS6* and *DL* act synergistically in terminating floral meristem.

RNA in situ hybridization analysis detected *DL* expression in the wild-type lemma and carpel, not in the palea (Yamaguchi et al., 2004) (see Supplemental Figures 1A to 1C online). However, ectopic expression of *DL* was observed in the altered palea organ and ectopic carpels or abnormal ovules in *mads6-1* (see Supplemental Figures 1D to 1G online) (Ohmori et al., 2009; Li et al., 2010). By contrast, *MADS6* expression was observed in the *dl-sup6* floral primordia at stage Sp4, and in palea and carpel primordia as well as the floral meristem at stages Sp7 and Sp8, just like in the wild type (see Supplemental Figures 1H to 1J online). Together, these results suggested that, whereas *MADS6* may repress the expression of *DL*, *DL* does not have an obvious effect on *MADS6* expression.

***MADS6* and *MADS13* Redundantly Regulate Carpel/Ovule Identity and Floral Determinacy**

MADS13 is a D-class gene that functions in specifying ovule identity and floral meristem determinacy (Dreni et al., 2007). We recently identified a strong *MADS13* allele, *mads13-3*, which showed carpelloid structures, indeterminate floral organs, and complete female sterility caused by aborted ovule development (Figures 5A, 5B, and 5K) (Li et al., 2011).

Flowers of the *mads6-1 mads13-3* double mutant displayed defects in the outer three whorls, a phenotype that is similar to that of *mads6-1* (Figures 5C and 5D). Given that the expression of *MADS13* is limited to ovules (Dreni et al., 2007) and that the *mads13* phenotype is exclusively found in the fourth whorl, we conclude that the function of *MADS6* in controlling the organ identities of the outer three whorls is independent of *MADS13*. However, the flowers of *mads6-1 mads13-3* displayed more severe defects in carpel/ovule development and floral meristem determinacy than the single mutants. We grouped these phenotypes into two categories: weak and strong. Fifty-seven percent ($n = 55$) of the double mutants had a weak phenotype, in which the ovule is transformed into carpelloid structures, a phenotype that is close to the severe phenotype shown in *mads13-3* or *mads6-1* (Figures 5F and 5I). Forty-three percent ($n = 55$) of the double mutants had the strong phenotype, in which floral meristem determinacy is largely lost due to the generation of higher order carpel-like organs along the floral axis (Figures 5E, 5J, and 5L). Consistent with these results was the detection of ectopic expression of the carpel marker gene *DL* in these ectopic carpel-like organs (see Supplemental Figures 1N and 1O online). Further scanning electron microscopy analysis showed that at stage Sp6, the phenotype of the *mads6-1 mads13-3* floral primordia was close to that of *mads6-1* (e.g., delayed stamen primordial development) (Figure 5G). However, after the carpel is formed at early Sp8 stage, the double mutant developed much stronger defects than the single mutants and had indeterminate floral meristems (Figure 5H). These results suggested that




Figure 4. Flower Phenotypes of *mads6-1 dl-sup6*.

(A) *dl-sup6* flower at stage Sp8.

(B) *dl-sup6* flower at stage Sp8, showing inner organs.

(C) and (D) *mads6-1 dl-sup6* flowers of the type I (C) and type II (D) phenotypes at stage Sp8. The arrow in (D) indicates the altered palea structure.

(E) *mads6-1 dl-sup6* flowers at stage Sp8, after the removal of the lemma and the palea. The arrow indicates one additional inflorescence-like structure in the center.

(F) to (H) Scanning electron microscopy analysis.

(F) *mads6-1 dl-sup6* flower at stage Sp8, showing an inflorescence-like structure in the center.

(G) *mads6-1 dl-sup6* flower at stage Sp6. Arrows indicate two palea-like structures.

(H) *mads6-1 dl-sup6* flower with inflorescence-like meristem at early stage Sp8.

(I) Transverse section of a *mads6-1 dl-sup6* flower at stage Sp8 showing two abnormal palea-like organs on the palea side.

(J) and (K) Floral diagrams of *dl-sup6* (J) and *mads6-1 dl-sup6* (K).

apa, abnormal palea; est, ectopic stamen; gl, glume-like organ; le, lemma; lo, lodicules; l-a, lodicule-anther mosaic organ; si, secondary inflorescence; st, stamen. Bars = 1 mm in (A) to (E), 500 μm in (F), 50 μm in (G) and (H), and 100 μm in (I).

MADS6 and *MADS13* have partial functional redundancy in specifying carpel/ovule identity and terminating floral stem cell activity.

In situ hybridization analysis detected the expression of *MADS13* in wild-type ovules at stage Sp8 (see Supplemental Figures 2A and 2B online), which is consistent with a previous report (Dreni et al., 2007). In *mads6-1*, *MADS13* expression was detected in ovule primordia at early Sp8 stage, similar to its expression in the wild type (see Supplemental Figure 2C online), and in ectopic ovules at late Sp8 stage (see Supplemental Figures 2D and 2E online). In *mads13-3*, the expression of *MADS6* was observed in the floral meristem and primordia of the palea, lodicules, and the carpel at stages Sp7 to Sp8 (see Supplemental Figures 2F to 2H online), which is similar to its

expression pattern in the wild type. These results suggested that *MADS6* and *MADS13* do not obviously regulate the expression of each other at the transcriptional level during early stages of ovule specification.

MADS6* Positively Regulates the Expression of *MADS7* and *MADS8

MADS7 and *MADS8* are the two closest rice homologs of *Arabidopsis SEP3* (Malcomber and Kellogg, 2005; Zahn et al., 2005; Arora et al., 2007). Like *SEP3*, *MADS7* and *MADS8* are expressed in the three inner whorls (Cui et al., 2010). *MADS6* was shown to interact with *MADS7* and *MADS8* in a yeast two-hybrid analysis (Moon et al., 1999). Moreover, Cui et al. (2010) revealed




Figure 5. Flower Phenotypes of *mads6-1 mads13-3*.

- (A) One *mads13-3* flower at stage Sp8 showing normal floral morphology.
 (B) Longitudinal section of one *mads13-3* carpel at stage Sp8 showing abnormal ovule.
 (C) One *mads6-1 mads13-3* flower at stage Sp8 displaying abnormal palea.
 (D) to (H) Scanning electron microscopy analysis.
 (D) One *mads6-1 mads13-3* flower at stage Sp8 displaying ectopic glume-like structures and abnormal lodicules and stamen development.
 (E) One *mads6-1 mads13-3* flower at stage Sp8 showing higher-order carpel-like structures in the center.
 (F) One *mads6-1 mads13-3* flower at stage Sp8 showing the transformation from ovule to carpel-like structure.
 (G) One *mads6-1 mads13-3* flower at stage Sp6 showing delayed inner floral organ development.
 (H) One *mads6-1 mads13-3* flower at early stage Sp8 showing new organ primordium in whorl 4.
 (I) and (J) Longitudinal sections of *mads6-1 mads13-3* flowers showing weak (I) and severe (J) phenotypes in the floral center.
 (K) and (L) Floral diagrams of *mads13-3* (K) and *mads6-1 mads13-3* (L).
 apa, abnormal palea; ca, carpel; ec, ectopic carpel; gl, glume-like organ; le, lemma; ll, lodicule-like organ; lo, lodicules; l-a, lodicule-anther mosaic organ; ov, ovule; pa, palea; st, stamen. Bars = 1 mm in (A), (C), and (D), 500 μ m in (E), 100 μ m in (B), (F), and (H) to (J), and 50 μ m in (G).

that plants in which both *MADS7* and *MADS8* were silenced displayed strong defects in flower development, where stamens were transformed into lodicules or glume-like organs and flowers appeared to lose determinacy by generating higher order carpels, suggesting functional redundancy between *MADS7* and *MADS8* in specifying flower development.

In situ hybridization detected the expression of *MADS7* in the primordia of lodicules, stamens, and carpels in the wild type (Figures 6A to 6C), which agrees with results from the previous report (Cui et al., 2010). In addition, the expression of *MADS8* was detected in the floral meristem (Figure 6D) and whorls 2, 3, and 4 (Figures 6E and 6F) in the wild type. In *mads6-1*, however, the expression of *MADS7* and *MADS8* was markedly reduced in floral meristem in the early stage of flower development (Figures 6G to 6L), which is consistent with the microarray data and qRT-PCR analysis (Table 1, Figure 1). Together, these re-

sults supported the notion that *MADS6* positively regulates the expression of *MADS7* and *MADS8* during rice flower development.

MADS6 Positively Regulates the Expression of *MADS3* and *MADS58*

In *mads6-1* flowers, the initiation of stamen development is much delayed and stamens are partially converted into lodicule-like or lodicule-anther mosaic organs during early flower development (Ohmori et al., 2009; Li et al., 2010). *MADS3*, a C-class gene in rice, has been shown to play a key role in controlling the development of lodicules and stamens (Yamaguchi et al., 2006; Hu et al., 2011). More recently we showed that *MADS3* also defines carpel development and floral determinacy redundantly with *MADS13* (Li et al., 2011). In addition, two intermediate




Figure 6. Expression Pattern of *MADS7* and *MADS8* in Wild-Type and *mads6-1* Flowers.

(A) Transcripts of *MADS7* in the primordia of the second and third whorls in the wild-type (WT) flower at early stage Sp6.

(B) and (C) *MADS7* expression signals in whorls 2, 3, and 4 of the wild-type flower at stage Sp7 (B) and Sp8 (C).

(D) to (F) *MADS8* expression in the wild-type flower at early stage Sp5 (D), Sp7 (E), and Sp8 (F).

(G) to (I) Expression of *MADS7* in *mads6-1* flowers at stage Sp6 (G), Sp7 (H), and Sp8 (I).

(J) to (L) Expression of *MADS8* in *mads6-1* flowers at stage Sp6 (J), Sp7 (K), and Sp8 (L).

le, lemma; pa, palea; lo, lodicules; st, stamen; ca, carpel. Arrows indicate lodicule primordia. Bars = 50 μm in (A), (D), (G), and (J), and 100 μm in (B), (C), (E), (F), (H), (I), (K), and (L).

alleles, *mads3-2* and *mads3-4*, displayed mild homeotic transformation from stamens into lodicule-like or lodicule-anther mosaic organs (Yamaguchi et al., 2006; Hu et al., 2011) (Figures 7A, 7F, and 7J).

The morphology of palea in the *mads6-1 mads3-4* double mutant appeared similar to *mads6-1*, suggesting that *MADS6*'s function in regulating palea identity is independent of *MADS3*. However, more severe defects were observed in whorls 2 to 4. In the double mutant, the average number of ectopic lodicule- or glume-like organs increased to 7.25 ($n = 91$), compared with 5.34 ($n = 79$) in *mads6-1*, and the number of stamens in the double mutant decreased to 2.45 ($n = 91$) from 6.00 in the wild type (Figures 7B, 7E, 7I, and 7K; see Supplemental Table 3 online). Defects in the double mutant also seemed to be intensified in whorl 4, where elongated floral axis or an inflorescence-like structure was observed in the floral center, and the average number of carpels was 2.2 ($n = 91$) (Figures 7C and 7D; see Supplemental Table 3 online). Scanning electron microscopy observation indicated that *mads6-1 mads3-4* had delayed inner floral organ development at stage Sp6 (Figure 7G), which is similar to that of *mads6-1*. At stage Sp8, the *mads6-1 mads3-4* double mutant displayed lodicule- or glume-like organs compared with the wild type or *mads3-4* (Figure 7H). These results suggested that *MADS6* and *MADS3* synergistically determine the identity of the three inner floral organs and floral determinacy.

In situ analysis detected the expression of *MADS3* at stage Sp5 in the wild type at the position where the stamen primordium was to be formed (Figure 8A). At stages Sp7 and Sp8, the mRNA of *MADS3* was detectable in whorls 3 and 4 of the wild-type

flowers (Figures 8B and 8C) (Yamaguchi et al., 2006). However, in *mads6-1*, the expression of *MADS3* seemed much weaker and delayed (Figures 8D to 8F). Similarly, reduced expression of *MADS58* was observed in *mads6-1* at the early stage (Figures 8G to 8J). This result is consistent with the microarray data and qRT-PCR analysis, suggesting that *MADS6* may promote the expression of *MADS3* and *MADS58* during early flower development. Conversely, we observed no obvious change in *MADS6* expression between *mads3-4* and the wild type, with the exception of some signals in ectopic lodicule-like primordia (Figures 8K to 8N), suggesting that *MADS3* does not affect the expression of *MADS6* at the transcriptional level to significant degrees.

The *mads58* allele contains a *dSpm* element insertion in the second intron of *MADS58*, which leads to ~ 35 -fold reduction in the expression of *MADS58* but no obvious defects in flower development (Figure 9A) (Dreni et al., 2011). The defects of floral organs in the outer three whorls were almost identical in *mads6-1 mads58* and *mads6-1*, except that the number of stigmas and abnormal carpels/ovules was higher in the double mutant (Figures 9B to 9G). The average stigma number per flower was 6.56 ($n = 50$) in *mads6-1 mads58* and 2.77 ($n = 44$) in *mads6-1*. About half of the *mads6-1 mads58* flowers displayed defective ovule development, each showing two or three carpel/ovule-like structures. The average number of carpel/ovule per flower was 1.5 ($n = 50$) in *mads6-1 mads58* and 1.1 in *mads6-1* ($n = 44$) under the same growth condition (Figures 9E and 9F).

MADS box proteins can regulate gene expression by binding as homo- or heterodimers to sequences containing a consensus




Figure 7. Flower Phenotypes of *mads6-1 mads3-4*.

- (A) *mads3-4* flower at stage Sp8. Lodicule-anther mosaic organs are indicated by arrows.
- (B) to (D) Flower phenotypes of *mads6-1 mads3-4* at stage Sp8, including many glume- or lodicule-like and lodicule-anther mosaic organs (B), two carpels in the center (C), and elongated flower axis or inflorescence-like organ and several carpel-like organs on the top (D).
- (E) to (H) Scanning electron microscopy analysis.
- (E) One *mads6-1 mads3-4* flower at stage Sp8 showing lodicule- and glume-like organs.
- (F) One *mads3-4* flower at Sp7 showing the primordium of one ectopic organ in the second whorl.
- (G) One *mads6-1 mads3-4* flower at stage Sp6 showing delayed development of inner floral organs, a similar phenotype to the *mads6-1* flowers.
- (H) One *mads6-1 mads3-4* flower at stage Sp8 showing primordia of lodicule- or glume-like organs.
- (I) Transverse section analysis of a stage-Sp8 *mads6-1 mads3-4* flower, which has six glume-like organs, four lodicule-like organs, two stamens, and two carpels.
- (J) and (K) Floral diagrams of *mads3-4* (J) and *mads6-1 mads3-4* (K).
- ca, carpel; ec, ectopic carpel; fm, floral meristem; gl, glume-like organ; l-a, lodicule-anther mosaic organ; ll, lodicules-like organ; lo, lodicules; st, stamen. Bars = 1 mm in (A) to (C), 500 μ m in (D) and (E), 50 μ m in (F) and (G), and 100 μ m in (H) and (I).

core element called the CArG box (Riechmann et al., 1996). For instance, the second intron of the *Arabidopsis* C-class gene *AG* contains one CArG-box, M1 [CC(AT)₆G], which is bound by SEP3 and acts as an enhancer sequence required for the proper expression of *AG* (Deyholos and Sieburth, 2000; Kaufmann et al., 2009) (Figure 10A). To further investigate the regulatory role of MADS6 during rice flower development, we searched the homeotic genes investigated in this study for putative CArG box sequences using the plant CARE tool (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>; Lescot et al., 2002). We found five CArG motifs in the second intron of *MADS58* and grouped them into two types: M2 [CC(AT)₈G] and M3 [CC(AT)₇G]. The second intron of *MADS3* was also found to contain five CArG motifs, which could be grouped into M2, M3, and M4 [CC(AT)₄(A/T/C/G)₂G] types (Figure 10A).

To test whether MADS6 is able to bind to the CArG motifs in the second introns of *MADS3* and *MADS58*, we performed chromatin immunoprecipitation (ChIP)-qPCR analysis. First, we developed rabbit polyclonal antibodies against a bacterial-expressed recombinant protein that contains 86 amino acids (amino acid 165 to amino acid 250) from the N terminus of MADS6 (see Methods). Specificity of the antibody was validated using immunoblot analysis, by which we detected, using a protein extract from wild-type flowers, a band of ~30 kD that is close to the expected size (see Supplemental Figure 3 online). ChIP-qPCR analysis using the affinity-purified MADS6 antibody showed specific enrichment of the M3-containing region of *MADS58* but not the other CArG motifs in *MADS3* and *MADS58* (Figure 10B; see Methods and Supplemental Table 4 online). These results suggested that MADS6 may directly regulate the expression of




Figure 8. Spatial and Temporal Expression Pattern of *MADS3*, *MADS58*, and *MADS6*.

(A) to (F) Expression pattern of *MADS3* in flowers of the wild type (WT) [(A) to (C)] and *mads6-1* [(D) to (F)]. (A) and (D), (B) and (E), and (C) and (F) are from stage Sp5, Sp7, and Sp8, respectively.

(G) to (J) Expression patterns of *MADS58* in flowers of the wild type [(G) and (H)] and *mads6-1* [(I) and (J)]. (G) and (I) and (H) and (J) are from stages 6 and 8, respectively.

(K) to (N) Expression pattern of *MADS6* in *mads3-4* flowers at stage Sp4 (K), Sp5 (L), and Sp6 [(M) and (N)]. The arrow in (N) indicates a lodicule-like or lodicule-anther mosaic organ.

ca, carpel; fm, floral meristem; gl, glume-like organ; lo, lodicule; pa, palea; st, stamen. Bars = 50 μm in (A), (D), (K), and (L) and 100 μm in (B), (C), (E) to (J), (M), and (N).

MADS58 through binding to the CArG element in the second intron.

DISCUSSION

MADS6 Is a Key Regulator in Specifying Floral Organ Identities

Flowering plants (angiosperms) evolved a tremendous diversity of floral structures (Theissen and Melzer, 2007). In this study, we investigated the genetic interaction of *MADS6* with floral homeotic genes *SPW1* (B-class), *MADS3* and *MADS58* (C-class), *MADS13* (D-class), and *DL* and demonstrated that interactions of *MADS6* with these floral homeotic genes play essential roles in rice flower development. Expression analyses indicated that the expression of B-, C-, and E-class genes in *mads6-1* was reduced at early flower developmental stages (Figure 1, Table 1). This may explain the delayed development of floral organ primordia in the mutant. However, repression of these six MADS box genes at the transcript level was less obvious at late flower developmental stages. Ohmori et al. (2009) reported the upregulation of *MADS14*, *MADS15*, *MADS3*, *MADS58*, and *LHS1* in *mfo1-1* elongated lodicules and *MADS14*, *MADS15*, and *MADS3* in the *mfo1-1* carpel during late flower developmental stages, suggesting the developmental stage-dependent control of these floral homeotic genes in rice. In this study, we did not detect obvious changes in the expression pattern of *MADS6* in *spw1-1*,

mads3-4, and *mads13-3*. Based on these results, we propose that *MADS6* may act as an upstream regulator that activates the expression of B- (*MADS4* and *SPW1*), C- (*MADS3* and *MADS58*), and E-class (*MADS7* and *MADS8*) genes during early flower development (Figure 11A). Alternatively, the reduced expression of B-, C-, and E-class genes may be a consequence of delayed development of flower organ primordia in *mads6-1*.

Phylogenetic and functional analyses revealed that *AGL6* family members are closely related to *SEP*-like genes (Becker and Theissen, 2003). *Arabidopsis SEP* and *AGL6* genes were shown to activate the expression of B- and C-class genes (Liu et al., 2009a; Koo et al., 2010). Moreover, ChIP sequencing analysis revealed direct association of *SEP3* with the regulatory regions of MADS box genes, where peaks of association were found to be located in the promoters of *AP1*, *AP3*, *SEP1*, and *SEP2* and the second intron of *AG* (Kaufmann et al., 2009). In this study, ChIP-qPCR assays revealed direct association of *MADS6* with a predicted regulatory motif in the second intron of *MADS58* (Figure 10). In addition, dramatically reduced expression of *MADS58* was observed in *mads6-1* during early flower development (Figures 1 and 8; Table 1), suggesting that *MADS6* functions upstream of the C-class gene *MADS58* and may directly activate *MADS58* expression (Figure 11A).

Similar to *SEP* proteins (Immink et al., 2009), *MADS6* have been shown to form complexes with *MADS4*, *SPW1*, *MADS13*, *MADS7*, *MADS8*, *MADS14*, and *MADS15* (Moon et al., 1999; Favaro et al., 2002; Seok et al., 2010) (Figure 11), suggesting that




Figure 9. Floral Phenotypes of *mads6-1 mads58*.

(A) One *mads58* flower at stage Sp8.

(B) to (D) *mads6-1 mads58* flowers at stage Sp8 showing the enlarged palea (B), inner floral organs (C), and an increased number of stigmas (D).

(E) and (F) Longitudinal sections of *mads6-1 mads58* flowers at stage Sp8 showing abnormal carpel/ovule development.

(G) Diagram of a *mads6-1 mads58* flower.

apa, abnormal palea; ca, carpel; gl, glume-like organs; le, lemma; ll, lodicule-like organs; lo, lodicules; l-a, lodicule-anther mosaic organs; ov, ovule; pa, palea; st, stamen; sti, stigmas. Bars = 1 mm in (A) to (C), 500 μ m in (D), and 100 μ m in (E) and (F).

AGL6-like proteins act as integrators that form multimeric complexes with MADS domain proteins from different clades in rice (Figure 11). We hypothesize that these MADS6-containing protein complexes may also be involved in the transcriptional control of floral homeotic genes in rice. This is exemplified by the findings that in *Arabidopsis* AP3 and PI form functional heterodimers, which can bind DNA *in vitro* (Riechmann et al., 1996) and regulate their own transcription *in vivo* (Goto and Meyerowitz, 1994; Krizek and Meyerowitz, 1996). Further determination of whether such auto/cross-regulatory loops also exist during flower development in grasses will help explain how *MADS6* acts both upstream from and in concert with these floral MADS box genes.

The ABCDE Model and Control of Palea Identity in Grasses

Compared with eudicot plants, rice has both conserved and unique genes that act in the regulation of flower development (Figure 11). As a result, the ABCDE genetic model is only partially applicable to the mechanism of flower development in grasses (Thompson and Hake, 2009; Li et al., 2011). For example, B- and C-class function seems to be conserved in grasses, *Arabidopsis*, and other eudicot species, whereas the function of *AGL6* genes, equivalents of the *Arabidopsis* class E or *SEP* genes, is partially conserved from eudicots to grasses (Figure 11).

In grasses, *AGL6*-like genes acquired a new expression domain in the palea, suggesting that their proteins may be key regulators of palea identity (Ohmori et al., 2009; Reinheimer and

Kellogg, 2009; Li et al., 2010). It has been hypothesized that during evolution of grass flowers, the palea was a congenital fusion of two perianth parts along with the lemma, which would represent a modified trimerous calyx (Francis, 1920; Zanis, 2007). Congenital fusion (also called phylogenetic fusion and zonal growth) refers to a compound structure that is developed as a homogeneous unit but is potentially derived from separate origins (Cusick, 1966; Verbeke, 1992). According to comparative morphological analysis, the development of compound organs resulting from fusions between individual primordia is thought to play a key role in floral morphogenesis during evolution (Verbeke, 1992). However, the mechanism underlying congenital fusion remains poorly understood.

Our results in this study suggest that both *MADS6* and *DL* are required for the proper establishment of palea identity and morphology. In the wild type, *DL* was shown to have zygomorphic expression in the lemma, but in *mads6* mutants, its expression becomes detectable in the palea as well. In addition, *mads6* mutants display loss of the palea identity (Ohmori et al., 2009; Li et al., 2010). *DL* was recently shown to play a critical role in specifying lemma identity (Figure 11), and *dl-sup6* lemmas had altered developmental identity with an increased number of vascular tissues but no obvious change of palea morphology (Li et al., 2011). Intriguingly, some flowers of the *mads6-1 dl-sup6* double mutants displayed complete loss of palea identity determination, leading to two separate organs. The complete loss of palea identity may be explained by the idea that congenital fusion




Figure 10. Analysis of the Association of MADS6 with the Predicted Regulatory Elements in the Three Homeotic Genes.

(A) Gene structures of *AG*, *MADS58*, and *MADS3* showing the conserved CArG motifs within their second introns.

(B) ChIP-PCR assay showing the specific enrichment of fragments that contain M3 from the second intron of *MADS58* by the MADS6 antibody. Error bars indicate SD ($n = 3$).

M2 to M5 represent motifs 2 to 5; IP, immunoprecipitated chromatin using MADS6 antibody; NoAb, negative control without MADS6 antibody.

[See online article for color version of this figure.]

of the palea relies on proper establishment of the palea identity. This interpretation is analogous to what was described as the dedoublement phenomenon in *Arabidopsis* stamens, where a single stamen primordium in the medial position is split in two, producing two pairs of medial stamens (Ronse Decraene and Smets, 1993; Bowman and Smyth, 1998). However, *ap3* or *pi* mutants show the transformation of these organs into carpels, resulting in just single organs in these positions (Bowman et al., 1991; Goto and Meyerowitz, 1994; Krizek and Meyerowitz, 1996). Similarly, this interpretation was used to explain the stamen number reduction in *Lepidium* (Brassicaceae) (i.e., stamens are reduced either by apparent loss of primordia [in the two lateral positions] or by fusion of two primordia into one [in the two medial positions]) (Endress, 1992; Bowman and Smyth, 1998). Here, we propose that MADS6 may specify the palea identity through repressing the expression of *DL* in the palea (Figure 11). Consistent with this view, investigations from model dicot plants suggest that the congenital fusion often acts downstream of organ identity specification such that a loss of organ identity alters proper fusion (Alvarez and Smyth, 1999; Vandebussche et al., 2004; Prunet et al., 2009). In *Arabidopsis*, analysis of double mutants of *crabs claw* (*crc*) and *spatula* (*spt*) with homeotic mutants indicated that A and B organ identity genes are capable of negatively regulating the function of *CRC* and *SPT* in carpel development (Alvarez and Smyth, 1999). *CRC* was shown to suppress the radial growth but trigger the longitudinal growth of the developing gynoecium, and *SPT* is able to promote the development of the carpel margins and the derived tissues (Alvarez and Smyth, 1999).

Previously, we showed that the rice *CYCLOIDEA*-like homolog *RETARDED PALEA1* (*REP1*) is a palea-specific gene that regulates palea identity and initiation by regulating cell proliferation and expansion, but it does not affect lemma development (Yuan et al., 2009). The *rep1* mutants display altered palea identity, such as an increased number of vascular bundles in the palea, which resembles a lemma-like organ (Yuan et al., 2009). Whether

and how MADS6 interacts with *REP1* in specifying palea identity requires further elucidation.

We have shown that in addition to specifying palea identity, *MADS6* also regulates lodicule development by interacting with *SPW1*, defines the stamen, carpel, and meristem identities with *MADS3* and *MADS58*, and specifies carpel/ovule development and floral meristem determinacy together with *MADS13* (Figure 11). Furthermore, *MADS6* regulates flower development redundantly with E-class genes, such as *LHS1*, *MADS7*, and *MADS8*. Although no obvious protein-protein interaction or transcriptional regulation between *MADS6* and *LHS1* was found (Li et al., 2010), *MADS6* may promote the expression of *MADS7* and *MADS8* and physically interact with the products of these two genes (Moon et al., 1999) (Figure 11). Similarly, *Arabidopsis* *AGL6* was shown to interact with several MADS proteins, such as *SEP1*, *SEP3*, *SHATTERPROOF2*, *AP1*, and *FUL* (de Folter et al., 2005).

Control of Floral Meristem Determinacy

Floral organs originate from the floral meristem, a pool of pluripotent stem cells (Liu et al., 2009b; Prunet et al., 2009). Mechanisms for the maintenance of floral meristem determinacy seem to be widely conserved among angiosperms (Ferrario et al., 2004; Prunet et al., 2009). Generally, floral meristem activity is abolished after the formation of a fixed pattern of floral organs. It has been shown that *Arabidopsis* *AG* is a key regulator in terminating floral meristem by turning *WUSCHEL* (*WUS*) off (Lenhard, et al., 2001; Lohmann et al., 2001). Whether grasses use a similar mechanism remains unknown. In the rice genome, there are 13 putative *WOX* (*WUSCHEL*-related homeobox gene family) genes; *Os WUS* was found to be closely related in sequence to the *Arabidopsis* *WUS* gene (Nardmann and Werr, 2006; Dai et al., 2007; Zhang et al., 2010b). But the biological function of *Os WUS* remains unclear. Unlike the *WUS* gene in *Arabidopsis*, *Os WUS* is not expressed in the organizing center of




Figure 11. Proposed Models for Rice Flower Development.

(A) Roles of MADS6 and floral homeotic genes in specifying rice flower development. *MADS6* specifies palea identity by repressing *DL* expression. It regulates lodicule development by activating the expression of *SPW1* and *MADS4* and physically interacting with the *MADS4*-*SPW1* complex (Seok et al., 2010). *MADS6* regulates the stamen, carpel, and meristem identities by activating the expression of *MADS3* and *MADS58* at early stages and specifies carpel/ovule development and controls floral meristem termination by interacting with *MADS13* (Moon et al., 1999). *MADS4* or *MADS13* interacts with *MADS7* and *MADS8*, respectively (Kater et al., 2006). In addition, *MADS6* determines flower development redundantly with *LHS1*, *MADS7*, and *MADS8* (Ohmori et al., 2009; Li et al., 2010). It also promotes the expression of *MADS7* and *MADS8* and physically interacts with *MADS7* and *MADS8* (Moon et al., 1999). Furthermore, *MADS6*, *MADS3*, *MADS58*, *DL*, and *MADS13* redundantly specify floral meristem determinacy (Yamaguchi et al., 2004, 2006; Dreni et al., 2011; Li et al., 2011).

(B) ABCDE model in rice flower development. The putative A-class gene *DEP* defines palea identity (Wang et al., 2010; Li et al., 2011). The E-function genes *LHS1*, *MADS6*, and *MADS34* specify lemma/palea identities. The A-class gene (*DEP*), B-class genes *SPW1*, *MADS2*, and *MADS4*, in combination with the E-class genes *LHS1*, *MADS6*, *MADS7*, *MADS8*, and *MADS34*, specify the lodicule identity. The B-class genes, the C-class genes *MADS3* and *MADS58*, together with the E-class genes determine the stamen identity. *DL* specifies carpel identity. The D-class gene *MADS13* and E-class genes determine carpel/ovule identity. Additionally, *DL* antagonistically regulates the expression of B-class genes, and the C-class gene *MADS3* represses the expression of the putative A-class gene *DEP* (*MADS15*) (Li et al., 2011).

ca, carpel; le, lemma; lo, lodicule; ov, ovule; pa, palea; st, stamen.

[See online article for color version of this figure.]

the vegetative shoot apical meristem (Nardmann and Werr, 2006). Grass species have duplicated C-class genes (Mena et al., 1996; Kramer et al., 2004; Yamaguchi et al., 2006; Zahn et al., 2006). In rice, analyses of *mads3* and *mads58* mutants suggested that *MADS3* plays a major role in specifying stamen identity and late anther development (Yamaguchi et al., 2006; Hu et al., 2011), and *MADS58* plays a redundant role with *MADS3* in specifying floral meristem determinacy and carpel identity (Dreni et al., 2011). More recently, our double mutant analysis revealed that *MADS3* and *MADS13* redundantly regulate carpel/

ovule development and floral meristem determinacy, suggesting that the C-class and D-class genes in rice retain their conserved function even though they had multiple subfunctionalization and/or neofunctionalization events after duplication within AG clade (Li et al., 2011) (Figure 11).

In this study, we have shown that *MADS6* regulates floral meristem determinacy together with C-class genes. Mutations in *MADS6* cause greatly reduced expression of the AG homologs *MADS3* and *MADS58*, and *mads6-1 mads3-4* displayed severe defects in floral meristem determinacy, such as the presence of inflorescence-like structures (Figure 7). Although *mads6-1 mads58* and *mads6-1* displayed similar defects in the outer three-whorl floral organs, the double mutant flowers had an increased number of stigma and carpel-like structures compared with *mads6-1* (Figure 9), indicating additional defects in floral determinacy. In maize, BDE and ZAG1 were shown to physically interact (Thompson et al., 2009), yet such an interaction was not found between *MADS6* and *MADS3/MADS58* (Moon et al., 1999). This may suggest that rice *MADS6* has conserved and divergent mechanisms from its counterpart in maize in forming protein complexes to specify floral development. Alternatively, *MADS6* may have weak interaction with *MADS3/58*, which was not detected by previous attempts (Moon et al., 1999).

Our data suggest that in addition to C-class genes, *MADS6* has other partners in determining floral meristem identity. *MADS6* functions redundantly with *SPW1*, *MADS13*, and *DL*, respectively, in specifying floral meristem identity. Some flowers in *mads6-1 spw1-1* exhibited new inflorescence-like organs in lieu of lodicules, suggesting that *MADS6* and *SPW1* play an important role in floral development, particularly in repressing meristem activity after the establishment of lodicule identity. Their genetic interaction may be explained by their protein-protein interaction observed by Seok et al. (2010). In addition to its role in promoting the expression of *MADS4* and *SPW1* during early flower development, *MADS6* was also shown to physically interact with *MADS4* and *SPW1*, suggesting that *MADS6*-*MADS4*-*SPW1* complex formation may be essential for flower development (Seok et al., 2010) (Figure 11). Additionally, analysis of *mads6-1 mad13-3* suggests the interaction of *MADS6* with *MADS13* in specifying floral meristem identity. *MADS6* can interact with *MADS13* at the protein level (Favaro et al., 2002), but they do not seem to regulate each other's expression as revealed in this study (Figure 11).

Rice *DL* is different from the well-known ABC genes (Nagasawa et al., 2003; Yamaguchi et al., 2004). Moreover, the role of *DL* is distinct from the closely related *Arabidopsis* *YABBY* gene *CRC*, which only plays a partial role in carpel identity (Alvarez and Smyth, 1999; Bowman and Smyth, 1999). Our recent analysis of *mads3-4 dl-sup6* flowers indicated that *DL* and *MADS3* act redundantly in specifying carpel identity and terminating floral meristem, but they may function in distinct pathways (Li et al., 2011) (Figure 11). Furthermore, previous investigation indicated that *DL* plays an antagonistic role with class B genes (Yamaguchi et al., 2004; Li et al., 2011). The *mads6-1 dl-sup6* double mutant flowers displayed inflorescence-like structures in whorl 4, suggesting that *MADS6* and *DL* may synergistically terminate the floral meristem (Figure 11).

MADS6 is able to specify floral meristem by interacting with E-class genes such as *LHS1/MADS1*, *MADS7*, and *MADS8* (Figure 11). One possible mechanism is that *MADS6* could positively regulate the expression of *MADS7* and *MADS8*. *mads1 mads6* double mutants had severe indeterminate floral meristem (Ohmori et al., 2009; Li et al., 2010), while our microarray (this study) and in situ analyses (Li et al., 2010) did not reveal obvious expression changes in *LHS1/MADS1* in *mads6-1*. Although Moon et al. (1999) reported the protein–protein interaction between *MADS6* and *LHS1/MADS1* in yeast cells, we did not observe this interaction (Li et al., 2010), possibly due to weak protein interaction.

In summary, using double mutant analyses in combination with in situ and ChIP-qPCR experiments, we reveal interactions of *MADS6* with floral homeotic genes *SPW1*, *MADS3*, *MADS58*, *MADS13*, and *DL* in regulating rice flower development and show that *MADS6* positively regulates the expression of the B-, C-, and E-class genes during early flower development. A model (Figure 11) is proposed to illustrate the role of floral homeotic genes in the specification of flower organ identity and meristem determination in rice.

METHODS

Plant Materials

The *mads6-1*, *mads13-3*, *mads3-4*, and *dl-sup6* mutants were identified previously (Chen et al., 2006b; Hu et al., 2011; Li et al., 2010, 2011). *spw1-1* and *mads58* were kindly provided by Hajime Sakai, Yasuo Nagato, and Venkatesan Sundaresan. Double mutants were isolated by phenotype observation and verified by genotyping. Methods for genotyping *dl-sup6*, *mads13-3*, and *mads3-4* were described previously (Li et al., 2011). Primer pairs 6TPF/6TPR, 16TPF/16TPR, and 58TPF/58TPR were used for genotyping *mads6-1*, *spw1-1*, and *mads58*, respectively (see Supplemental Table 4 online). Mutant and wild-type rice (*Oryza sativa*) plants were planted in the paddy field or greenhouse in Shanghai Jiao Tong University, China.

Histological Analysis and Microscopy Observation

Flower materials were collected at various stages of development according to spikelet size and morphology definitions described by Ikeda et al. (2004), fixed in FAA (10% formalin, 50% ethanol, and 5% acetic acid), and dehydrated in a series of graded ethanol (Chu et al., 2006; Li et al., 2006). For histological analysis, tissues were then infiltrated with xylene and embedded in paraplast plus. Then, materials were sectioned to 8 μm thick and stained with toluidine blue (Bio Basic) and photographed using a Nikon E600 microscope and a Nikon DXM1200 digital camera. Scanning electron microscopy was performed with a JSM-6360LV (Jeol) as described previously (Li et al., 2006).

In Situ Hybridization

Samples were treated as described previously (Li et al., 2006). Constructs of gene-specific probes for *MADS13*, *SPW1*, *DL*, and *MADS6* were described previously (Li et al., 2010, 2011). Constructs of *MADS3* and *MADS58* probes were generated based on descriptions by Yamaguchi et al. (2006) and probes of *MADS7* and *MADS8* were made as described by Cui et al. (2010). Digoxigenin-labeled antisense and sense probes were transcribed in vitro as described by Li et al. (2006). Images were obtained using an Olympus Nikon E600 microscope.

Microarray Analysis

Microarray experiments were performed as described by Hu et al. (2011). Total RNAs were isolated from three replicates of stage Sp6 flowers from the wild type and *mads6-1*. Developing flower primordia were collected based on microscopy analysis and spikelet length described by Ikeda et al. (2004).

qRT-PCR

Total RNA was isolated from wild-type and *mads6-1* flowers at stage Sp4/SP5, Sp6, early Sp8, and late Sp8. qRT-PCR conditions were the same as those described by Zhang et al. (2010a). Primers for *MADS3* and *MADS4* were described by Chen et al. (2006a), and primers for *SPW1*, *MADS7*, *MADS8*, and *MADS58* were described by Cui et al. (2010). All samples were run with three replicates. Data acquisition and analyses were performed using the Roche Light Cycler software. Sample amounts used were normalized using *ACTIN* expression.

Preparation of the *MADS6* Polyclonal Antibody

The *MADS6*-specific fragment (493 to 753 bp) was amplified by PCR using the primer pair 6APF/6APR (see Supplemental Table 4 online). PCR products were cloned into pET-32a (Novagen) to produce p32-P-*MADS6*. The 6XHis-*MADS6* fused protein was expressed in *Escherichia coli* using p32-P-*MADS6*. Proteins were purified according to the manufacturer's instructions, and the antibody was prepared as described by Huang et al. (2003) using the specific *MADS6* fragment. Nuclear extracts from the wild-type flowers at stage Sp7 were used for immunoblot analysis to test the specificity of the *MADS6* antibody, following the protocol used for the ChIP experiments (Zhang et al., 2010b), except that the tissue was not fixed. The glutathione S-transferase-tagged full-length *MADS6* protein was used as the positive control for immunoblot analysis.

ChIP-qPCR and Fold Enrichment Analysis

Rice spikelets at stage Sp7 were treated and sonicated with an Ultrasonic Crasher Noise Isolating Chamber (Scientz). The procedure for ChIP of the *MADS6*-DNA complexes in wild-type flowers was modified from Haring et al. (2007). For each PCR reaction, 0.5 μL of recovered DNA from immune precipitation (IP) or mock was used as template, three biological replicates were included, and each reaction was repeated three times. Primers used for qPCR analyses are labeled in Figure 10 and listed in Supplemental Table 4 online, and reactions were performed on a Rotor-Gene RG3000A detection system (Corbett Research) using SYBR Green I. The normalized mean cycle threshold (Ct) of each gene was calculated and used to determine fold change according to the method described by Rotor-Gene version 6.0 (Build 38) software and Zhang et al. (2010a). The difference between the Ct of the antibody enrichment and no antibody control (mock) was calculated to obtain the relative enrichment of the fragments containing the putative CARG motifs. Quantification involved normalization of the Ct of each IP or the control sample to obtain a nonspecific adjustment ΔCt ($\Delta\text{Ct IP} - \Delta\text{Ct mock}$), followed by calculation of relative enrichment of each fragment using the equation $2^{-(\Delta\text{Ct IP} - \Delta\text{Ct control})}$.

Accession Numbers

Sequence data from this article for the cDNAs of *MADS6*, *SPW1*, *MADS3*, *MADS58*, *DL*, *MADS13*, *MADS7*, and *MADS8* can be found in the GenBank/EMBL data libraries under accession numbers AK069103, AK069317, AK108568, AK111723, AK242416, AK070425, AK100263, and AK072867, respectively. Locus identifications in the Rice Genome

Annotation Project Database are as follows: *MADS6* (Os02g45770), *SPW1* (Os06g49840), *MADS3* (Os01g10504), *MADS58* (Os05g11414), *DL* (Os03g11600), *MADS13* (Os12g10540), *MADS7* (Os08g41950), and *MADS8* (Os09g32948). Microarray data accession number in NCBI is GSE29349.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure 1. Spatial and Temporal Expression Pattern of *DL* and *MADS6* as Revealed by in Situ Hybridization.

Supplemental Figure 2. Spatial and Temporal Expression Pattern of *MADS13* and *MADS6*.

Supplemental Figure 3. Specificity Analysis of *MADS6* Polyclonal Antibodies.

Supplemental Table 1. Up- or Downregulated (at Least Twofold Change in Expression, P Value < 0.05) Non-MADS Box Genes in *mads6-1* Identified by Bayes Analysis (<0.5% FDR).

Supplemental Table 2. Number of Floral Organs in Wild-Type and *mads6-1*, *spw1-1*, and *mads6-1 spw1-1* Flowers.

Supplemental Table 3. Number of Floral Organs in Wild-Type and *mads6-1*, *mads3-4*, and *mads6-1 mads3-4* Flowers.

Supplemental Table 4. Primers Used in This Study.

ACKNOWLEDGMENTS

We thank Zhijing Luo and Mingjiao Chen for mutant screening and generation of mutants, Hajime Sakai and Yasuo Nagato for providing *spw1-1*, Venkatesan Sundaresan for *mads58*, Ning Jiang and Zhongchi Liu for comments on the this work, and Jianping Hu for editing this manuscript. We gratefully acknowledge the anonymous reviewers for very helpful comments. This work was supported by Funds from the National Basic Research Program of China (2009CB941500 and 2007CB108700), the National Natural Science Foundation of China (30725022), the Chinese Transgenic Project (2009ZX08009-108B), and the National 863 High-Tech Project (2011AA10A101).

AUTHOR CONTRIBUTIONS

H.L., W.L., Y.H., L.Z., C.Y., J.X., L.D., and M.M.K. performed research and analyzed data. D.Z. and W.L. designed the research and analyzed data. D.Z. wrote the article.

Received May 10, 2011; revised June 21, 2011; accepted July 5, 2011; published July 22, 2011.

REFERENCES

- Agrawal, G.K., Abe, K., Yamazaki, M., Miyao, A., and Hirochika, H. (2005). Conservation of the E-function for floral organ identity in rice revealed by the analysis of tissue culture-induced loss-of-function mutants of the *OsMADS1* gene. *Plant Mol. Biol.* **59**: 125–135.
- Alvarez, J., and Smyth, D.R. (1999). *CRABS CLAW* and *SPATULA*, two *Arabidopsis* genes that control carpel development in parallel with *AGAMOUS*. *Development* **126**: 2377–2386.
- Ambrose, B.A., Lerner, D.R., Ciceri, P., Padilla, C.M., Yanofsky, M.F., and Schmidt, R.J. (2000). Molecular and genetic analyses of the *silky1* gene reveal conservation in floral organ specification between eudicots and monocots. *Mol. Cell* **5**: 569–579.
- Arora, R., Agarwal, P., Ray, S., Singh, A.K., Singh, V.P., Tyagi, A.K., and Kapoor, S. (2007). MADS-box gene family in rice: Genome-wide identification, organization and expression profiling during reproductive development and stress. *BMC Genomics* **8**: 242.
- Becker, A., and Theissen, G. (2003). The major clades of MADS-box genes and their role in the development and evolution of flowering plants. *Mol. Phylogenet. Evol.* **29**: 464–489.
- Bowman, J.L., and Smyth, D.R. (1998). Patterns of petal and stamen reduction in reduction in Australian species of *Lepidium* L. (Brassicaceae). *Int. J. Plant Sci.* **159**: 65–74.
- Bowman, J.L., and Smyth, D.R. (1999). *CRABS CLAW*, a gene that regulates carpel and nectary development in *Arabidopsis*, encodes a novel protein with zinc finger and helix-loop-helix domains. *Development* **126**: 2387–2396.
- Bowman, J.L., Smyth, D.R., and Meyerowitz, E.M. (1991). Genetic interactions among floral homeotic genes of *Arabidopsis*. *Development* **112**: 1–20.
- Chen, L., Chu, H.W., Yuan, Z., Pan, A.H., Liang, W.Q., Huang, H., Shen, M.S., Zhang, D.B., and Chen, L. (2006b). Isolation and genetic analysis for rice mutants treated with ⁶⁰Co γ -Ray. *J. Xiamen Univ.* **45**: 82–85.
- Chen, Z.X., Wu, J.G., Ding, W.N., Chen, H.M., Wu, P., and Shi, C.H. (2006a). Morphogenesis and molecular basis on *naked seed rice*, a novel homeotic mutation of *OsMADS1* regulating transcript level of *AP3* homologue in rice. *Planta* **223**: 882–890.
- Chu, H.W., Qian, Q., Liang, W.Q., Yin, C.S., Tan, H.X., Yao, X., Yuan, Z., Yang, J., Huang, H., Luo, D., Ma, H., and Zhang, D.B. (2006). The *floral organ number4* gene encoding a putative ortholog of *Arabidopsis CLAVATA3* regulates apical meristem size in rice. *Plant Physiol.* **142**: 1039–1052.
- Clayton, W.D., and Renvoize, S.A. (1986). *Genera Graminum: Grasses of the World*. (Kew, UK: Kew Publishing).
- Clifford, H. (1987). *Spikelet and Floral Morphology*. (Washington, DC: Smithsonian Institution Press).
- Coen, E.S., and Meyerowitz, E.M. (1991). The war of the whorls: Genetic interactions controlling flower development. *Nature* **353**: 31–37.
- Colombo, L., Franken, J., Koetje, E., van Went, J., Dons, H.J., Angenot, G.C., and van Tunen, A.J. (1995). The petunia MADS box gene *FBP11* determines ovule identity. *Plant Cell* **7**: 1859–1868.
- Cui, R., Han, J., Zhao, S., Su, K., Wu, F., Du, X., Xu, Q., Chong, K., Theissen, G., and Meng, Z. (2010). Functional conservation and diversification of class E floral homeotic genes in rice (*Oryza sativa*). *Plant J.* **61**: 767–781.
- Cusick, F. (1966). On phylogenetic and ontogenetic fusions. In *Trends in Plant Morphogenesis*, E.G. Cutter, ed (London: Longmans, Green & Co.), pp. 170–183.
- Dai, M., Hu, Y., Zhao, Y., Liu, H., and Zhou, D.X. (2007). A *WUSCHEL-LIKE HOMEBOX* gene represses a *YABBY* gene expression required for rice leaf development. *Plant Physiol.* **144**: 380–390.
- de Folter, S., Immink, R.G., Kieffer, M., Parenicová, L., Henz, S.R., Weigel, D., Busscher, M., Kooiker, M., Colombo, L., Kater, M.M., Davies, B., and Angenot, G.C. (2005). Comprehensive interaction map of the *Arabidopsis* MADS Box transcription factors. *Plant Cell* **17**: 1424–1433.
- Deyholos, M.K., and Sieburth, L.E. (2000). Separable whorl-specific expression and negative regulation by enhancer elements within the *AGAMOUS* second intron. *Plant Cell* **12**: 1799–1810.
- Ditta, G., Pinyopich, A., Robles, P., Pelaz, S., and Yanofsky, M.F. (2004). The *SEP4* gene of *Arabidopsis thaliana* functions in floral organ and meristem identity. *Curr. Biol.* **14**: 1935–1940.

- Dreni, L., Jacchia, S., Fornara, F., Fornari, M., Ouwerkerk, P.B., An, G., Colombo, L., and Kater, M.M. (2007). The D-lineage MADS-box gene *OsMADS13* controls ovule identity in rice. *Plant J.* **52**: 690–699.
- Dreni, L., Pilatone, A., Yun, D.P., Erreni, S., Pajoro, A., Caporali, E., Zhang, D.B., and Kater, M.M. (2011). Functional analysis of all AGAMOUS subfamily members in rice reveals their roles in reproductive organ identity determination and meristem determinacy. *The Plant Cell*, in press.
- Endress, P.K. (1992). Evolution and floral diversity: The phylogenetic surroundings of *Arabidopsis* and *Antirrhinum*. *Int. J. Plant Sci.* **153** (suppl.): S106–S122.
- Favaro, R., Immink, R.G., Ferioli, V., Bernasconi, B., Byzova, M., Angenent, G.C., Kater, M., and Colombo, L. (2002). Ovule-specific MADS-box proteins have conserved protein-protein interactions in monocot and dicot plants. *Mol. Genet. Genomics* **268**: 152–159.
- Ferrario, S., Immink, R.G., and Angenent, G.C. (2004). Conservation and diversity in flower land. *Curr. Opin. Plant Biol.* **7**: 84–91.
- Francis, M.E. (1920). *Book of Grasses*. (Garden City, New York: Doubleday, Page & Co.).
- Gao, X., Liang, W., Yin, C., Ji, S., Wang, H., Su, X., Guo, C., Kong, H., Xue, H., and Zhang, D. (2010). The *SEPALLATA*-like gene *OsMADS34* is required for rice inflorescence and spikelet development. *Plant Physiol.* **153**: 728–740.
- Goto, K., and Meyerowitz, E.M. (1994). Function and regulation of the *Arabidopsis* floral homeotic gene *PISTILLATA*. *Genes Dev.* **8**: 1548–1560.
- Grass Phylogeny Working Group (2001). Phylogeny and subfamilial classification of the grasses (Poaceae). *Ann. Mo. Bot. Gard.* **88**: 373–457.
- Haring, M., Offermann, S., Danker, T., Horst, I., Peterhansel, C., and Stam, M. (2007). Chromatin immunoprecipitation: Optimization, quantitative analysis and data normalization. *Plant Methods* **3**: 11.
- Hu, L.F., Liang, W.Q., Yin, C.S., Cui, X., Zong, J., Wang, X., Hu, J.P., and Zhang, D.B. (2011). Rice *MADS3* regulates ROS homeostasis during late anther development. *Plant Cell* **23**: 515–533.
- Huang, Y., Liang, W., Pan, A., Zhou, Z., Huang, C., Chen, J., and Zhang, D. (2003). Production of FaeG, the major subunit of K88 fimbriae, in transgenic tobacco plants and its immunogenicity in mice. *Infect. Immun.* **71**: 5436–5439.
- Ikeda, K., Nagasawa, N., and Nagato, Y. (2004). Developmental course of inflorescence and spikelet in rice. *Breed. Sci.* **54**: 147–156.
- Immink, R.G., Tonaco, I.A., de Folter, S., Shchennikova, A., van Dijk, A.D., Busscher-Lange, J., Borst, J.W., and Angenent, G.C. (2009). *SEPALLATA3*: The ‘glue’ for MADS box transcription factor complex formation. *Genome Biol.* **10**: R24.
- Jeon, J.S., Jang, S., Lee, S., Nam, J., Kim, C., Lee, S.H., Chung, Y.Y., Kim, S.R., Lee, Y.H., Cho, Y.G., and An, G. (2000). *leafy hull sterile1* is a homeotic mutation in a rice MADS box gene affecting rice flower development. *Plant Cell* **12**: 871–884.
- Kater, M.M., Dreni, L., and Colombo, L. (2006). Functional conservation of MADS-box factors controlling floral organ identity in rice and *Arabidopsis*. *J. Exp. Bot.* **57**: 3433–3444.
- Kaufmann, K., Muiño, J.M., Jauregui, R., Airoidi, C.A., Smaczniak, C., Krajewski, P., and Angenent, G.C. (2009). Target genes of the MADS transcription factor *SEPALLATA3*: Integration of developmental and hormonal pathways in the *Arabidopsis* flower. *PLoS Biol.* **7**: e1000090.
- Kobayashi, K., Maekawa, M., Miyao, A., Hirochika, H., and Kyojuka, J. (2010). *PANICLE PHYTOMER2 (PAP2)*, encoding a *SEPALLATA* subfamily MADS-box protein, positively controls spikelet meristem identity in rice. *Plant Cell Physiol.* **51**: 47–57.
- Koo, S.C., et al. (2010). Control of lateral organ development and flowering time by the *Arabidopsis thaliana* MADS-box Gene *AGAMOUS-LIKE6*. *Plant J.* **62**: 807–816.
- Kramer, E.M., Jaramillo, M.A., and Di Stilio, V.S. (2004). Patterns of gene duplication and functional evolution during the diversification of the *AGAMOUS* subfamily of MADS box genes in angiosperms. *Genetics* **166**: 1011–1023.
- Krizek, B.A., and Meyerowitz, E.M. (1996). The *Arabidopsis* homeotic genes *APETALA3* and *PISTILLATA* are sufficient to provide the B class organ identity function. *Development* **122**: 11–22.
- Kyojuka, J., Kobayashi, T., Morita, M., and Shimamoto, K. (2000). Spatially and temporally regulated expression of rice MADS box genes with similarity to *Arabidopsis* class A, B and C genes. *Plant Cell Physiol.* **41**: 710–718.
- Lenhard, M., Bohnert, A., Jürgens, G., and Laux, T. (2001). Termination of stem cell maintenance in *Arabidopsis* floral meristems by interactions between *WUSCHEL* and *AGAMOUS*. *Cell* **105**: 805–814.
- Lescot, M., Déhais, P., Thijs, G., Marchal, K., Moreau, Y., Van de Peer, Y., Rouzé, P., and Rombauts, S. (2002). PlantCARE, a database of plant *cis*-acting regulatory elements and a portal to tools for *in silico* analysis of promoter sequences. *Nucleic Acids Res.* **30**: 325–327.
- Li, H.F., Liang, W.Q., Jia, R.D., Yin, C.S., Zong, J., Kong, H.Z., and Zhang, D.B. (2010). The *AGL6*-like gene *OsMADS6* regulates floral organ and meristem identities in rice. *Cell Res.* **20**: 299–313.
- Li, H.F., Liang, W.Q., Yin, C.S., Zhu, L., and Zhang, D.B. (2011). Genetic interaction of *OsMADS3*, *DROOPING LEAF*, and *OsMADS13* in specifying rice floral organ identities and meristem determinacy. *Plant Physiol.* **156**: 263–274.
- Li, N., et al. (2006). The rice *tapetum degeneration retardation* gene is required for tapetum degradation and anther development. *Plant Cell* **18**: 2999–3014.
- Linder, H., and Rudall, P. (2005). Evolutionary history of poales. *Annu. Rev. Ecol. Evol. Syst.* **36**: 107–124.
- Liu, C., Thong, Z., and Yu, H. (2009b). Coming into bloom: The specification of floral meristems. *Development* **136**: 3379–3391.
- Liu, C., Xi, W.Y., Shen, L.S., Tan, C.P., and Yu, H. (2009a). Regulation of floral patterning by flowering time genes. *Dev. Cell* **16**: 711–722.
- Lohmann, J.U., Hong, R.L., Hobe, M., Busch, M.A., Parcy, F., Simon, R., and Weigel, D. (2001). A molecular link between stem cell regulation and floral patterning in *Arabidopsis*. *Cell* **105**: 793–803.
- Malcomber, S.T., and Kellogg, E.A. (2004). Heterogeneous expression patterns and separate roles of the *SEPALLATA* gene *LEAFY HULL STERILE1* in grasses. *Plant Cell* **16**: 1692–1706.
- Malcomber, S.T., and Kellogg, E.A. (2005). *SEPALLATA* gene diversification: Brave new whorls. *Trends Plant Sci.* **10**: 427–435.
- Mena, M., Ambrose, B.A., Meeley, R.B., Briggs, S.P., Yanofsky, M.F., and Schmidt, R.J. (1996). Diversification of C-function activity in maize flower development. *Science* **274**: 1537–1540.
- Moon, Y.H., Kang, H.G., Jung, J.Y., Jeon, J.S., Sung, S.K., and An, G. (1999). Determination of the motif responsible for interaction between the rice *APETALA1/AGAMOUS-LIKE9* family proteins using a yeast two-hybrid system. *Plant Physiol.* **120**: 1193–1204.
- Münster, T., Deleu, W., Wingen, L.U., Ouzunova, M., Cacharrón, J., Faigl, W., Werth, S., Kim, J.T.T., Saedler, H., and Theissen, G. (2002). Maize MADS-box genes galore. *Maydica* **47**: 287–301.
- Nagasawa, N., Miyoshi, M., Sano, Y., Satoh, H., Hirano, H., Sakai, H., and Nagato, Y. (2003). *SUPERWOMAN1* and *DROOPING LEAF* genes control floral organ identity in rice. *Development* **130**: 705–718.
- Nardmann, J., and Werr, W. (2006). The shoot stem cell niche in angiosperms: Expression patterns of *WUS* orthologues in rice and maize imply major modifications in the course of mono- and dicot evolution. *Mol. Biol. Evol.* **23**: 2492–2504.
- Ohmori, S., Kimizu, M., Sugita, M., Miyao, A., Hirochika, H., Uchida, E., Nagato, Y., and Yoshida, H. (2009). *MOSAIC FLORAL ORGANS1*, an *AGL6*-like MADS box gene, regulates floral organ identity and meristem fate in rice. *Plant Cell* **21**: 3008–3025.

- Pelaz, S., Ditta, G.S., Baumann, E., Wisman, E., and Yanofsky, M.F. (2000). B and C floral organ identity functions require *SEPALLATA* MADS-box genes. *Nature* **405**: 200–203.
- Pinyopich, A., Ditta, G.S., Savidge, B., Liljegen, S.J., Baumann, E., Wisman, E., and Yanofsky, M.F. (2003). Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature* **424**: 85–88.
- Prasad, K., Parameswaran, S., and Vijayraghavan, U. (2005). Os-MADS1, a rice MADS-box factor, controls differentiation of specific cell types in the lemma and palea and is an early-acting regulator of inner floral organs. *Plant J.* **43**: 915–928.
- Preston, J.C., and Kellogg, E.A. (2006). Reconstructing the evolutionary history of paralogous *APETALA1/FRUITFULL*-like genes in grasses (Poaceae). *Genetics* **174**: 421–437.
- Prunet, N., Morel, P., Negrutiu, I., and Trehin, C. (2009). Time to stop: Flower meristem termination. *Plant Physiol.* **150**: 1764–1772.
- Prusinkiewicz, P., Erasmus, Y., Lane, B., Harder, L.D., and Coen, E. (2007). Evolution and development of inflorescence architectures. *Science* **316**: 1452–1456.
- Reinheimer, R., and Kellogg, E.A. (2009). Evolution of *AGL6*-like MADS box genes in grasses (Poaceae): Ovule expression is ancient and palea expression is new. *Plant Cell* **21**: 2591–2605.
- Riechmann, J.L., Wang, M., and Meyerowitz, E.M. (1996). DNA-binding properties of Arabidopsis MADS domain homeotic proteins *APETALA1*, *APETALA3*, *PISTILLATA* and *AGAMOUS*. *Nucleic Acids Res.* **24**: 3134–3141.
- Rijkema, A.S., Zethof, J., Gerats, T., and Vandenbussche, M. (2009). The petunia *AGL6* gene has a *SEPALLATA*-like function in floral patterning. *Plant J.* **60**: 1–9.
- Ronse Decraene, L.P., and Smets, E.F. (1993). Dedoublement revisited: Towards a renewed interpretation of the androecium of the Magnoliophytina. *Bot. J. Linn. Soc.* **113**: 103–124.
- Rudall, P.J., Stuppy, W., Cunniff, J., Kellogg, E.A., and Briggs, B.G. (2005). Evolution of reproductive structures in grasses (Poaceae) inferred by sister-group comparison with their putative closest living relatives, *Ecdeiocoleaceae*. *Am. J. Bot.* **92**: 1432–1443.
- Schauer, S.E., Baskar, R., Brand, L., Bolanos, A., Grobei, M., Federer, M., Jauch, U., and Gagliardini, V. (2007). Examination of the role of the Arabidopsis MADS-box transcription factors *AGL6* and *AGL13* in reproduction. *Dev. Biol.* **306**: 312.
- Seok, H.Y., Park, H.Y., Park, J.I., Lee, Y.M., Lee, S.Y., An, G., and Moon, Y.H. (2010). Rice ternary MADS protein complexes containing class B MADS heterodimer. *Biochem. Biophys. Res. Commun.* **401**: 598–604.
- Smyth, G.K. (2004). Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3**: Article3.
- Theissen, G. (2001). Development of floral organ identity: stories from the MADS house. *Curr. Opin. Plant Biol.* **4**: 75–85.
- Theissen, G., Becker, A., Di Rosa, A., Kanno, A., Kim, J.T., Münster, T., Winter, K.U., and Saedler, H. (2000). A short history of MADS-box genes in plants. *Plant Mol. Biol.* **42**: 115–149.
- Theissen, G., and Melzer, R. (2007). Molecular mechanisms underlying origin and diversification of the angiosperm flower. *Ann. Bot. (Lond.)* **100**: 603–619.
- Theissen, G., and Saedler, H. (2001). Plant biology. Floral quartets. *Nature* **409**: 469–471.
- Thompson, B.E., Bartling, L., Whipple, C., Hall, D.H., Sakai, H., Schmidt, R., and Hake, S. (2009). *bearded-ear* encodes a MADS box transcription factor critical for maize floral development. *Plant Cell* **21**: 2578–2590.
- Thompson, B.E., and Hake, S. (2009). Translational biology: From *Arabidopsis* flowers to grass inflorescence architecture. *Plant Physiol.* **149**: 38–45.
- Vandenbussche, M., Theissen, G., Van de Peer, Y., and Gerats, T. (2003a). Structural diversification and neo-functionalization during floral MADS-box gene evolution by C-terminal frameshift mutations. *Nucleic Acids Res.* **31**: 4401–4409.
- Vandenbussche, M., Zethof, J., Royaert, S., Weterings, K., and Gerats, T. (2004). The duplicated B-class heterodimer model: Whorl-specific effects and complex genetic interactions in *Petunia hybrida* flower development. *Plant Cell* **16**: 741–754.
- Vandenbussche, M., Zethof, J., Souer, E., Koes, R., Tornielli, G.B., Pezzotti, M., Ferrario, S., Angenent, G.C., and Gerats, T. (2003b). Toward the analysis of the petunia MADS box gene family by reverse and forward transposon insertion mutagenesis approaches: B, C, and D floral organ identity functions require *SEPALLATA*-like MADS box genes in petunia. *Plant Cell* **15**: 2680–2693.
- Verbeke, J.A. (1992). Fusion events during floral morphogenesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **43**: 583–598.
- Wang, K., Tang, D., Hong, L., Xu, W., Huang, J., Li, M., Gu, M., Xue, Y., and Cheng, Z. (2010). *DEP* and *AFO* regulate reproductive habit in rice. *PLoS Genet.* **6**: e1000818.
- Whipple, C.J., Ciceri, P., Padilla, C.M., Ambrose, B.A., Bandong, S.L., and Schmidt, R.J. (2004). Conservation of B-class floral homeotic gene function between maize and *Arabidopsis*. *Development* **131**: 6083–6091.
- Whipple, C.J., Zanis, M.J., Kellogg, E.A., and Schmidt, R.J. (2007). Conservation of B class gene expression in the second whorl of a basal grass and outgroups links the origin of lodicules and petals. *Proc. Natl. Acad. Sci. USA* **104**: 1081–1086.
- Yamaguchi, T., Lee, D.Y., Miyao, A., Hirochika, H., An, G., and Hirano, H.Y. (2006). Functional diversification of the two C-class MADS box genes *OSMADS3* and *OSMADS58* in *Oryza sativa*. *Plant Cell* **18**: 15–28.
- Yamaguchi, T., Nagasawa, N., Kawasaki, S., Matsuoka, M., Nagato, Y., and Hirano, H.Y. (2004). The *YABBY* gene *DROOPING LEAF* regulates carpel specification and midrib development in *Oryza sativa*. *Plant Cell* **16**: 500–509.
- Yoo, S.K., Wu, X.L., Lee, J.S., and Ahn, J.H. (2011). *AGAMOUS-LIKE 6* is a floral promoter that negatively regulates the FLC/MAF clade genes and positively regulates FT in Arabidopsis. *Plant J.* **65**: 62–76.
- Yuan, Z., Gao, S., Xue, D.W., Luo, D., Li, L.T., Ding, S.Y., Yao, X., Wilson, Z.A., Qian, Q., and Zhang, D.B. (2009). *RETARDED PALEA1* controls palea development and floral zygomorphy in rice. *Plant Physiol.* **149**: 235–244.
- Zahn, L.M., Kong, H., Leebens-Mack, J.H., Kim, S., Soltis, P.S., Landherr, L.L., Soltis, D.E., Depamphilis, C.W., and Ma, H. (2005). The evolution of the *SEPALLATA* subfamily of MADS-box genes: A preangiosperm origin with multiple duplications throughout angiosperm history. *Genetics* **169**: 2209–2223.
- Zahn, L.M., Leebens-Mack, J.H., Arrington, J.M., Hu, Y., Landherr, L.L., dePamphilis, C.W., Becker, A., Theissen, G., and Ma, H. (2006). Conservation and divergence in the *AGAMOUS* subfamily of MADS-box genes: evidence of independent sub- and neofunctionalization events. *Evol. Dev.* **8**: 30–45.
- Zanis, M.J. (2007). Grass spikelet genetics and duplicate gene comparisons. *Int. J. Plant Sci.* **168**: 93–110.
- Zhang, D.B., and Wilson, Z.A. (2009). Stamen specification and anther development in rice. *Chin. Sci. Bull.* **54**: 1–12.
- Zhang, H., Liang, W.Q., Yang, X.J., Luo, X., Jiang, N., Ma, H., and Zhang, D.B. (2010a). *Carbon starved anther* encodes a MYB domain protein that regulates sugar partitioning required for rice pollen development. *Plant Cell* **22**: 672–689.
- Zhang, X., Zong, J., Liu, J., Yin, J.Y., and Zhang, D.B. (2010b). Genome-wide analysis of *WOX* gene family in rice, sorghum, maize, Arabidopsis and poplar. *J. Integr. Plant Biol.* **52**: 1016–1026.